



US007244585B1

(12) **United States Patent**  
**Sangar et al.**

(10) **Patent No.:** **US 7,244,585 B1**  
(45) **Date of Patent:** **Jul. 17, 2007**

(54) **3' SEQUENCE OF THE GB VIRUS B (GBV-B) GENOME**

(75) Inventors: **David V. Sangar**, Burbage Leics (GB);  
**Stanley M. Lemon**, Galveston, TX (US)

(73) Assignee: **The Board of Regents of The University of Texas System**, Austin, TX (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1230 days.

(21) Appl. No.: **09/587,653**

(22) Filed: **Jun. 5, 2000**

**Related U.S. Application Data**

(60) Provisional application No. 60/137,665, filed on Jun. 4, 1999.

(51) **Int. Cl.**

- C12P 19/36* (2006.01)
- C12N 15/09* (2006.01)
- C12N 15/51* (2006.01)
- C12N 7/01* (2006.01)
- C07H 19/00* (2006.01)

(52) **U.S. Cl.** ..... **435/69.1**; 435/320.1; 435/91.4; 424/228.1; 536/24.1

(58) **Field of Classification Search** ..... 435/235.1, 435/237, 239, 325, 5, 69.1; 536/23.72, 23.7; 424/204.1, 225.1

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

- 5,672,350 A \* 9/1997 Parker et al. .... 424/221.1
- 5,807,670 A 9/1998 Muerhoff et al. .... 435/5
- 6,627,437 B1 \* 9/2003 Traboni ..... 435/320.1

**OTHER PUBLICATIONS**

- Buckwold et al. *Antiviral Research* 2005, vol. 66, pp. 165-168.\*
- Bukh et al. *Virology* 1999, vol. 262, No. 2, pp. 470-478.\*
- Sbardellati et al. *J. Gene. Virol.* 2001, vol. 82, pp. 2437-2448.\*
- NCBI AJ277947 p. 1-5.\*
- The HIV Life Cycle, published by OpenChemist.Net on Jun. 5, 2006, pp. 1-15.\*
- Brass et al. *Int. J. Med.* 2006, vol. 3, pp. 29-34.\*
- Bartenschlager et al. *J. Gene. Viro.* 2000, vol. 81, pp. 1631-1648.\*
- Virus Replication published by Microbiology @ Leicester: *Virology* on Oct. 22, 2004, pp. 1-20.\*
- The Free Dictionary by FARLEX on line, Google searches on Aug. 11, 2006, pp. 1-2.\*
- Al et al., "Expression of recombinant hepatitis C virus in non-structural protein 5B *Escherichia coli*," *Virus Res* 53:141-149, 1998.
- Beard et al. "An infectious molecular clone of a Japanese genotype 1b hepatitis C virus," *Hepatology*, 30:316-324 1999.
- Behrens et al., "Identification and properties of the RNA dependent RNA polymerase of hepatitis C virus," *EMBO J*, 15:12-22, 1996.

Chambers et al., "Yellow fever/Japanese encephalitis chimeric viruses. construction and biological properties," *J Virol*, 73:3095-3101, 1999.

Choo et al., "Vaccination of chimpanzees against infection by the hepatitis C virus," *Proc Natl Acad Sci USA*, 91:1294-1298, 1994.

Farci et al., "Lack of protective immunity against reinfection with hepatitis C virus," *Science*, 258:135-140, 1992.

Farci et al., "The natural history of infection with hepatitis C virus (HCV) in chimpanzees: Comparison of serologic responses measured with first- and second-generation assays and relationship to HCV viremia," *The Journal of Infectious Diseases*, 165:1006-1011, 1992.

Frolov et al., "Cis-acting RNA elements required for replication of bovine viral diarrhea virus-hepatitis C virus 5' nontranslated region chimeras," *RNA* 4:1418-1435, 1998.

Hijikata et al., "Two distinct protease activities required for the processing of a putative nonstructural precursor protein of hepatitis C virus," *J Virol*, 67:4665-4675, 1993.

Honda et al., "Stability of a stem-loop involving the initiator AUG controls the efficiency of internal initiation of translation on hepatitis C virus RNA," *RNA*, 2:955-968, 1996.

Karayiannis et al., "Studies of GB hepatitis agent in tamarins," *Hepatology*, 9:186-192, 1989.

Kim et al., "Crystal structure of the hepatitis C virus NS3 protease domain complexed with a synthetic NS4A cofactor peptide," *Cell*, 87:343-355, 1996.

Kolykhalov et al., "Transmission of hepatitis C by intrahepatic inoculation of transcribed RNA," *Science*, 277:570-574, 1997.

Kolykhalov et al., "Identification of a highly conserved element at the 3' terminus of hepatitis C virus genome RNA," *J Virol*, 70:3363-3371, 1996.

Lemon and Honda, "Internal ribosome entry sites within the RNA genomes of hepatitis C virus and other flaviviruses," *Seminars in Virology*, 8:274-288, 1997.

Lu and Wimmer, "Poliovirus chimeras replicating under the translational control of genetic elements of hepatitis C virus reveal unusual properties of the internal ribosomal entry site of hepatitis C virus," *Proc Natl Acad Sci USA*, 93:1412-1417, 1996.

Muerhoff et al., "Genomic organization of GB viruses A and B: Two new members of the flaviviridae associated with GB agent hepatitis," *J Virol*, 69:5621-5630, 1995.

Pletnev et al., "Construction and characterization of chimeric tick-borne encephalitis/dengue type 4 viruses," *Proc. Natl Acad Sci USA*, 89:10532-10536, 1992.

(Continued)

*Primary Examiner*—Bruce R. Campell  
*Assistant Examiner*—Qun Li Bao  
(74) *Attorney, Agent, or Firm*—Fulbright & Jaworski L.L.P.

(57) **ABSTRACT**

The present invention relates generally to the fields of biochemistry, molecular biology, and virology. More particularly, it relates to the identification of 259 nucleotides located at the 3' end of the GB virus B (GBV-B) genome. The invention involves nucleic acid constructs and compositions encoding GBV-B sequence, including the 3' end of the sequence, which has allowed an infectious GBV-B clone to be constructed. This construct, and chimeric versions of it, may be employed to study GBV-B and related hepatitis family members, such as hepatitis C virus. The invention thus includes methods of preparing GBV-B-containing sequences, constructs, and viruses, as well as methods of employing these compositions.

OTHER PUBLICATIONS

Reynolds et al., "Unique features of internal initiation of hepatitis C virus RNA translation," *EMBO J*, 14:6010-6020, 1995.

Rijnbrand et al., "Mutational analysis of the GB virus B internal ribosome entry site," *J Virol*, 74:773-783, 2000.

Scarselli et al., "GB virus B and hepatitis C virus share substrate specificity," *J Virol*, 74:4985-4989, 1997.

Schlauder et al., "Molecular and serological analysis in the transmission of the GB hepatitis agent," *J Med Virol*, 46:81-90, 1995.

Shaffer et al., "A hepatitis A virus deletion mutant which lacks the first pyrimidine-rich tract of the 5' nontranslated RNA remains virulent in primates after direct intrahepatic nucleic acid transfection," *J Virol*, 69:6600-6604, 1995.

Simons et al., "Identification of two flavivirus like genomes in the GB hepatitis agent," *Proc Natl Acad Sci*, 92:3401-3405, 1995.

Tanaka et al., "A novel sequence found at the 3' terminus of hepatitis C virus genome," *Biochem Biophys Res Comm*, 215:744-749, 1995.

Todd et al., "Replication competent picornaviruses with complete genomic RNA 3' non-coding deletions," *J Virol*, 71:8868-8874, 1997.

Yanagi et al., "Transcripts from a single full length cDNA clone of hepatitis C virus are infectious when directly transfected into a liver of a chimpanzee," *Proc Natl Acad Sci USA*, 94:8738-8743, 1998.

Yao et al., "Structure of the hepatitis C virus RNA helicase domain," *Nature Structural Biology*, 4:463-467, 1997.

Zhao et al., "Poliovirus/hepatitis C virus (internal ribosome entry site-core) chimeric viruses: improved growth properties through modification of a proteolytic cleavage site and requirement for core RNA sequence but not for core related polypeptides," *J Virol*, 73:1546-1554, 1999.

\* cited by examiner




FIG. 1

### 3' SEQUENCE OF THE GB VIRUS B (GBV-B) GENOME

The present application claims the benefit of U.S. Provisional Application Ser. No. 60/137,665 filed on Jun. 4, 1999. The entire text of the above-referenced disclosure is herein incorporated by reference.

#### BACKGROUND OF THE INVENTION

##### A. Field of the Invention

The present invention relates generally to the fields of biochemistry, molecular biology, and virology. More particularly, it relates to the identification of 259 nucleotides of previously unrecognized sequence located at the 3' end of the GB virus B (GBV-B) genome.

##### B. Description of Related Art

Chronic hepatitis C is a major threat to the public health. Serologic surveys suggest that as many as 3.9 million Americans are chronically infected with the responsible virus, hepatitis C virus (HCV) (Alter, 1997). These individuals are at increased risk of developing progressive hepatic fibrosis leading to cirrhosis and loss of hepatocellular function, as well as hepatocellular carcinoma. The course of chronic hepatitis C is typically lengthy, often extending over decades, with insidious clinical progression usually occurring in the absence of symptoms. Nonetheless, liver disease due to HCV results in the death of 8,000-10,000 Americans annually, and chronic hepatitis C is the most common cause of liver transplantation within the U.S.

Therefore, HCV is a major public health problem. However, therapy for chronic hepatitis C is problematic. Recombinant interferon- $\alpha$  is approved for treatment of chronic hepatitis C (Consensus Development Panel, 1997). The benefit of interferon- $\alpha$  results primarily from its antiviral properties and its ability to inhibit production of virus by infected hepatocytes (Neumann et al., 1998). Nonetheless, even under optimal therapeutic regimens, the majority of patients with chronic hepatitis C fail to eliminate the virus or resolve their liver disease. Treatment failures are especially common in persons infected with genotype 1 HCV, unfortunately the most prevalent genotype in the U.S. Thus, there is an urgent need to better understand the virus and develop better treatment. Unfortunately, technical difficulties in working with HCV have made it necessary to use infectious surrogate viruses in efforts to develop treatments and vaccines for HCV.

Scientists' efforts to better understand HCV and to develop new drugs for treatment of hepatitis C have been stymied by two overwhelming technical deficiencies: first, the nonexistence of a high permissive cell line that supports replication of the virus and second, the absence of a permissive animal species other than chimpanzees, which are endangered and therefore available on a limited basis.

Presently, those who are working on HCV treatment and prevention are employing an infectious chimeric virus of sindbis and HCV and/or an infectious clone of pestivirus as surrogate virus models in HCV drug discovery efforts, due to the above technical difficulties of working with HCV. Alternatively, they are using isolated proteins or RNA segments of HCV for biochemical and structural studies. This approach precludes functional studies of virus replication and its inhibition.

GBV-B is a hepatotropic flavivirus that has a unique phylogenetic relationship to human HCV and strong potential to serve as a surrogate virus in drug discovery efforts related to hepatitis C antiviral drug development. GBV-B

causes acute hepatitis in experimentally infected tamarins (Simons et al., 1995; Schlauder et al., 1995; Karayiannis et al., 1989) and can serve as a surrogate virus for HCV in drug discovery efforts (due to technical difficulties in working with HCV). GBV-B virus is much closer in sequence and biological properties than the above-described models. It will be easier to make biologically relevant chimeras between HCV and GBV-B than by using more distantly related viruses. GBV-B is hepatotropic (as is HCV), whereas the viruses used in these competing technologies are not. In view of the above, an infectious clone of GBV-B would be useful to those working on HCV treatment and prevention.

Unfortunately, the use of GBV-B as a surrogate or model for HCV has not been possible in the past, because no infectious molecular clone of GBV-B virus genome could be prepared. It is now known that this obstacle was encountered because the GBV-B genome was believed to be 259 nucleotides shorter than its actual length (Muerhoff et al., 1995; Simons et al., 1995). Others, previous to the inventors, had failed to realize that the 3' sequence of GBV-B was missing from the prior sequences. Without this 3' sequence, it is not possible to prepare an infectious GBV-B molecular clone.

#### SUMMARY OF THE INVENTION

As discussed above, an infectious molecular clone of GBV-B would be very useful for the development of HCV preventative and therapeutic treatments. The construction of an infectious molecular clone of this virus will require the newly determined 3' sequence to be included in order for the clone to be viable. The inventors have elucidated the previously unrecognized 3' terminal sequence of GBV-B (SEQ ID NO:1). This sequence has been reproducibly recovered from tamarin serum containing GBV-B RNA, in RT-PCR protocols using several different primer sets, and as a fusion with previously reported 5' GBV-B sequences.

The newly identified 3' sequence is not included in published reports of the GBV-B sequence, nor described in patents relating to the original identification of the viral sequence (see U.S. Pat. No. 5,807,670 and references therein).

The invention has utility in that the inclusion of the sequence will be necessary for construction of an infectious molecular GBV-B clone. Such clones clearly have the potential to be constructed as chimeras including relevant hepatitis C virus sequences in lieu of the homologous GBV-B sequence, providing unique tools for drug discovery efforts. A full-length molecular clone of GBV-B was constructed, as described in later sections of this specification.

GBV-B can be used as a model for HCV, and the GBV-B genome can be used as the acceptor molecule in the construction of chimeric viral RNAs containing sequences of both HCV and GBV-B. Such studies will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (e.g., proteinase) required for HCV replication. However, all this work is dependent upon construction of an infectious clone of GBV-B, which is itself dependent on the incorporation of the correct 3' terminal nucleotide sequence within this clone. GBV-B has unique advantages over HCV in terms of its ability to replicate and cause liver disease in tamarins, which present fewer restrictions to research than chimpanzees, the only nonhuman primate species known to be permissive for HCV.

An infectious molecular clone of GBV-B is expected to have utility in liver-specific gene expression or in gene

therapy. This application might be enhanced by the inclusion of HCV genomic sequence in the form of a GBV-B/HCV chimera. Further, an infectious GBV-B/HCV chimera expressing HCV envelope proteins can have utility as a vaccine immunogen for hepatitis C.

A full-length cDNA copy of the GBV-B genome was constructed to contain the newly identified 3' terminal sequences. RNA transcribed from this cDNA copy of the genome would be infectious when inoculated into the liver of a GBV-B permissive tamarin, giving rise to rescued GBV-B virus particles. A chimeric molecule would then be constructed from this infectious GBV-B clone in which the HCV NS3 proteinase or proteinase/helicase sequence (or other relevant HCV sequences of interest in drug discovery efforts) would be placed in frame in lieu of the homologous GBV-B sequence, and this chimeric cDNA would be used to generate infectious GBV-B/HCV chimeric viruses by intrahepatic inoculation of synthetic RNA in tamarins. Published studies indicate that the GBV-B and HCV proteinases have closely related substrate recognition and cleavage properties, making such chimeras highly likely to be viable. These newly generated chimeric GBV-B/HCV viruses could be used in preclinical testing of candidate HCV NS3 proteinase inhibitors.

Therefore, the present invention encompasses an isolated polynucleotide encoding a 3' sequence of the GBV-B genome. The polynucleotide may include the sequence identified as SEQ ID NO:1. It is contemplated that the polynucleotide may be a DNA molecule or it can RNA molecule. It is further contemplated that expression constructs may contain a polynucleotide that has a stretch of contiguous nucleotides from SEQ ID NO:1 and/or SEQ ID NO:2, for example, lengths of 50, 100, 150, 250, 500, 1000, 5000, as well as the entire length of SEQ ID NO:1 or 2, are considered appropriate. Such polynucleotides may also be contained in other constructs of the invention or be used in the methods of the invention. Polynucleotides employing sequences from SEQ ID NO:1 may alternatively contain sequences from SEQ ID NO:2 in the constructs and methods of the present invention.

The invention is also understood as covering a viral expression construct that includes a polynucleotide encoding a 3' sequence of the GBV-B genome. This expression construct is further understood to contain the sequence identified as SEQ ID NO:1. The present invention contemplates the expression construct as a plasmid or as a virus. Furthermore, the expression construct can express GBV-B sequences; alternatively it may express sequences from a chimeric GBV-B/HCV virus.

The identification and isolation of a 3' sequence of GBV-B additionally provides a method of producing a virus, particularly a full-length virus, by introducing into a host cell a viral expression construct containing a polynucleotide encoding a 3' sequence of GBV-B and by culturing the host cell under conditions permitting production of a virus from the construct. This method can be practiced using a prokaryotic cell as a host cell, or by using a eukaryotic cell as a host cell. Furthermore, the eukaryotic cell can be located within an animal.

A method of producing virus according to the claimed invention can also be employed using a polynucleotide that contains synthetic RNA and/or synthetic DNA. Moreover, a step can be added to the method by also isolating any virus produced from the host cell. The virus can then be purified to homogeneity.

In further embodiments, the present invention encompasses an oligonucleotide between about 10 and about 259

consecutive bases of SEQ ID NO:1. This oligonucleotide is contemplated to be about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 50 bases in length, about 100 bases in length, about 150 bases in length, about 200 bases in length, or about 259 bases in length.

Additional examples of the claimed invention include a method for identifying a compound active against a viral infection by providing a virus expressed from a viral construct containing a 3' sequence of a GBV-B virus, by contacting the virus with a candidate substance; and by comparing the infectious ability of the virus in the presence of the candidate substance with the infectious ability of the virus in a similar system in the absence of the candidate substance. It is contemplated that the invention can be practiced using GBV-B virus or a GBV-B/HCV chimera.

The present invention can also be understood to provide a compound active against a viral infection identified by providing a virus expressed from a viral construct containing a 3' sequence of a GBV-B virus; contacting the virus with a candidate substance; and comparing the infectious ability of the virus in the presence of the candidate substance with the infectious ability of the virus in a similar system in the absence of the candidate substance. In some embodiments an active compound is identified using a GBV-B virus, while in other embodiments an active compound is identified using a GBV-B/HCV chimera.

As used herein the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" may mean one or more than one. As used herein "another" may mean at least a second or more.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1. Predicted secondary RNA structure at the 3' end of the novel 3' GBV-B sequence. The structure shown is that predicted for the 3' terminal 108 nucleotides of the GBV-B genomic RNA by the MFOLD 3.0 computer program of Zucker and Turner, and has an initial free energy of -48 kcal/mole. The predicted structure contains 3 stem-loops, numbered from the 3' end of the genome and labeled SL-1, SL-2, and SL-3. The structure of SL-1 is highly probable, given its terminal position within the genome. Alternative foldings of SL-2 and SL-3 are possible.

## 5

DESCRIPTION OF ILLUSTRATIVE  
EMBODIMENTS

## A. GVB-B Virus

The GBV-B genome structure is very similar to hepatitis C and these viruses share approximately 25% nucleotide identity (Simons et al., 1995; Muerhoff et al., 1995). As indicated above, this makes GBV-B more closely related to HCV than any other known virus. GBV-B genomic RNA is about 9.5 kb in length (Muerhoff et al., 1995) with a structured 5' noncoding region that contains an IRES that shares many structural features with the HCV IRES (Honda et al., 1996; Rijnbrand et al., 1999). As in HCV, this IRES drives the cap-independent translation of a long open reading frame. The polyprotein expressed from this reading frame appears to be organized identically to that of HCV, and processed to generate proteins with functions similar to those of HCV (Muerhoff et al., 1995). In fact, the major serine proteinases of these viruses (NS3) have been shown to have similar cleavage specificities (Scarselli et al., 1997). Finally, like HCV and distinct from the pestiviruses, the genomic RNA of GBV-B has a poly(U) tract located near its 3' terminus (Simons et al., 1995; Muerhoff et al., 1995). In addition, unreported sequences located at the extreme 3' end of the genome have been identified. This work indicates that the GBV-B RNA, like that of HCV (HCV (Tanaka et al., 1995; Kolykhalov et al., 1996), terminates in a lengthy run of heterogeneous bases (310 nts in GBV-B) possessing a readily apparent secondary structure

## B. NUCLEIC ACIDS

The present invention provides a nucleic acid sequence encoding a 3' sequence of the GBV-B genome (SEQ ID NO:1).

It should be clear that the present invention is not limited to the specific nucleic acids disclosed herein. As discussed below, a "3' sequence of the GBV-B genome" may contain a variety of different bases and yet still be functionally indistinguishable from the sequences disclosed herein. Such functionally indistinguishable sequences are likely to maintain the basic structure depicted in FIGS. 1 and 2, which may be used to guide the prediction of viable nucleotide substitutions.

## 1. Polynucleotides Encoding the 3' Sequence of the GBV-B Genome

A 3' sequence of the GBV-B genome disclosed in SEQ ID NO:1 is one aspect of the present invention. Nucleic acids according to the present invention may encode the 3' sequence of the GBV-B genome set forth in SEQ ID NO:1, the entire GBV-B genome, or any other fragment of a 3' sequence of the GBV-B genome set forth herein. The nucleic acid may be derived from genomic RNA as cDNA, i.e., cloned directly from the genome of GBV-B. cDNA may also be assembled from synthetic oligonucleotide segments.

It also is contemplated that a 3' sequence of the GBV-B genome may be represented by natural variants that have slightly different nucleic acid sequences but, nonetheless, maintain the same general structure (see FIGS. 1 and 2) and perform the same function in RNA replication.

As used in this application, the term "a nucleic acid encoding a 3' sequence of the GBV-B genome" refers to a nucleic acid molecule that may be isolated free of total viral nucleic acid. In preferred embodiments, the invention concerns nucleic acid sequences essentially as set forth in SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ

## 6

ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ID NO:12. The term "as set forth in SEQ ID NO:1" means that the nucleic acid sequence substantially corresponds to a portion of SEQ ID NO:1. It is contemplated that the techniques and methods described in this disclosure may apply to any of the sequences contained herein, including SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, and SEQ ED NO:12.

Allowing for the degeneracy of the genetic code, sequences that have at least about 50%, usually at least about 60%, more usually about 70%, most usually about 80%, preferably at least about 90% and most preferably about 95% of nucleotides that are identical to the nucleotides of SEQ ID NO:1 will be sequences that are "as set forth in SEQ ID NO:1." Sequences that are essentially the same as those set forth in SEQ ID NO:1 may also be functionally defined as sequences that are capable of hybridizing to a nucleic acid segment containing the complement of SEQ ID NO:1 under standard conditions.

The nucleic acid segments and polynucleotides of the present invention include those encoding biologically functional equivalent 3' sequences of the GBV-B genome. Changes designed by man may be introduced through the application of site-directed mutagenesis techniques or may be introduced randomly and screened later for the desired function, as described below.

3' sequence of the GBV-B genome sequences also are provided. Each of the foregoing is included within all aspects of the following description. The present invention concerns cDNA segments reverse transcribed from GBV-B genomic RNA (referred to as "DNA"). As used herein, the term "polynucleotide" refers to an RNA or DNA molecule that may be isolated free of other RNA or DNA of a particular species.

"Isolated substantially away from other coding sequences" means that the 3' sequence of the GBV-B genome forms the significant part of the RNA or DNA segment and that the segment does not contain large portions of naturally-occurring coding RNA or DNA, such as large fragments or other functional genes or cDNA noncoding regions. Of course, this refers to the polynucleotide as originally isolated, and does not exclude genes or coding regions later added to the it by the hand of man.

In certain other embodiments, the invention concerns isolated DNA segments (cDNA segments reverse transcribed from GVB-B genomic RNA) and recombinant vectors that include within their sequence a nucleic acid sequence essentially as set forth in SEQ ID NO:1. The term "essentially as set forth in SEQ ID NO:1" is used in the same sense as described above.

It also will be understood that nucleic acid sequences may include additional residues, such as additional 5' or 3' sequences, and still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological activity. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include additional various non-coding sequences flanking either of the 5' or 3' portions of the coding region, which are known to occur within viral genomes.

Sequences that are essentially the same as those set forth in SEQ ID NO:1 also may be functionally defined as sequences that are capable of hybridizing to a nucleic acid segment containing the complement of SEQ ID NO:1 under



For example, a medium stringency condition could be provided by about 0.1 to 0.25 M NaCl at temperatures of about 37° C. to about 55° C., while a low stringency condition could be provided by about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20° C. to about 55° C. Thus, hybridization conditions can be readily manipulated and thus will generally be a method of choice depending on the desired results.

In other embodiments, hybridization may be achieved under conditions of, for example, 50 mM Tris-HCl (pH 8.3), 75 mM KCl, 3 mM MgCl<sub>2</sub>, 10 mM dithiothreitol, at temperatures between approximately 20° C. to about 37° C. Other hybridization conditions utilized could include approximately 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, at temperatures ranging from approximately 40° C. to about 72° C. Formamide and SDS also may be used to alter the hybridization conditions.

One method of using probes and primers of the present invention is in the search for other viral sequences related to GBV-B or, more particularly, homologs of the GBV-B sequence. By varying the stringency of hybridization, and the region of the probe, different degrees of homology may be discovered.

Another way of exploiting probes and primers of the present invention is in site-directed, or site-specific, mutagenesis. The technique provides a ready ability to prepare and test sequence variants, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into complementary DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences that encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered.

The technique typically employs a bacteriophage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage vectors are commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis, which eliminates the step of transferring the gene of interest from a phage to a plasmid.

In general, site-directed mutagenesis is performed by first obtaining a single-stranded vector, or melting of two strands of a double stranded vector which includes within its sequence a DNA sequence encoding the desired protein. An oligonucleotide primer bearing the desired mutated sequence is synthetically prepared. This primer is then annealed with the single-stranded DNA preparation, taking into account the degree of mismatch when selecting hybridization conditions, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected that include recombinant vectors bearing the mutated sequence arrangement. There are newer and simpler site-directed mutagenesis techniques that can also be employed for this purpose. These include

procedures marketed in kit form that are readily available to one of ordinary skill in the art.

The preparation of sequence variants of the selected gene using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be limiting, as there are other ways in which sequence variants of genes may be obtained. For example, recombinant vectors encoding the desired gene may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

### 3. Antisense Constructs

In certain embodiments of the invention, the use of antisense constructs of the 3' sequence of the GBV-B genome is contemplated.

Antisense methodology takes advantage of the fact that nucleic acids tend to pair with "complementary" sequences. By complementary, it is meant that polynucleotides are those which are capable of base-pairing according to the standard Watson-Crick complementarity rules. That is, the larger purines will base pair with the smaller pyrimidines to form combinations of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T) in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. Inclusion of less common bases such as inosine, 5-methylcytosine, 6-methyladenine, hypoxanthine and others in hybridizing sequences does not interfere with pairing.

Targeting double-stranded (ds) DNA with polynucleotides leads to triple-helix formation; targeting RNA will lead to double-helix formation. Antisense polynucleotides, when introduced into a target cell, specifically bind to their target polynucleotide and interfere with transcription, RNA processing, transport, translation and/or stability. Antisense RNA constructs, or DNA encoding such antisense RNAs, may be employed to inhibit gene transcription or translation or both within a host cell, either in vitro or in vivo, such as within a host animal, including a human subject.

Antisense constructs could be used to block early steps in the replication of GBV-B and related viruses, by annealing to 3' terminal sequences and blocking their role in negative-strand initiation.

As stated above, "complementary" or "antisense" means polynucleotide sequences that are substantially complementary over their entire length and have very few base mismatches. For example, sequences of 15 bases in length may be termed complementary when they have complementary nucleotides at 13 or 14 positions. Naturally, sequences which are completely complementary will be sequences which are entirely complementary throughout their entire length and have no base mismatches. Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology but also contains a non-homologous region (e.g., ribozyme; see below) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

### 4. Amplification and PCR™

The present invention utilizes amplification techniques in a number of its embodiments. Nucleic acids used as a template for amplification are isolated from cells contained in the biological sample, according to standard methodologies (Sambrook et al., 1989). The nucleic acid may be genomic DNA or RNA or fractionated or whole cell RNA. Where RNA is used, it may be desired to convert the RNA to a complementary DNA using reverse transcriptase (RT). In one embodiment, the RNA is genomic RNA and is used directly as the template for amplification. In others, genomic



RNA is first converted to a complementary DNA sequence (cDNA) and this product is amplified according to protocols described below.

Pairs of primers that selectively hybridize to nucleic acids corresponding to GBV-B sequences are contacted with the isolated nucleic acid under conditions that permit selective hybridization. The term "primer," as defined herein, is meant to encompass any nucleic acid that is capable of priming the synthesis of a nascent nucleic acid in a template-dependent process. Typically, primers are oligonucleotides from ten to twenty base pairs in length, but longer sequences can be employed. Primers may be provided in double-stranded or single-stranded form, although the single-stranded form is preferred.

Once hybridized, the nucleic acid:primer complex is contacted with one or more enzymes that facilitate template-dependent nucleic acid synthesis. Multiple rounds of amplification, also referred to as "cycles," are conducted until a sufficient amount of amplification product is produced.

Next, the amplification product is detected. In certain applications, the detection may be performed by visual means. Alternatively, the detection may involve indirect identification of the product via chemiluminescence, radioactive scintigraphy of incorporated radiolabel or fluorescent label or even via a system using electrical or thermal impulse signals.

A number of template dependent processes are available to amplify the marker sequences present in a given template sample. One of the best known amplification methods is the polymerase chain reaction (referred to as PCR<sup>TM</sup>) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, and each incorporated herein by reference in entirety.

Briefly, in PCR<sup>TM</sup>, two or more primer sequences are prepared that are complementary to regions on opposite complementary strands of the marker sequence. An excess of deoxynucleoside triphosphates are added to a reaction mixture along with a DNA polymerase, e.g., Taq polymerase. If the marker sequence is present in a sample, the primers will bind to the marker and the polymerase will cause the primers to be extended along the marker sequence by adding on nucleotides. By raising and lowering the temperature of the reaction mixture, the extended primers will dissociate from the marker to form reaction products, excess primers will bind to the marker and to the reaction products and the process is repeated.

A reverse transcriptase PCR<sup>TM</sup> amplification procedure may be performed in order to quantify the amount of mRNA amplified. Methods of reverse transcribing RNA into cDNA are well known and described in Sambrook et al., 1989. Alternative methods for reverse transcription utilize thermostable, RNA-dependent DNA polymerases. These methods are described in WO 90/07641, filed Dec. 21, 1990, incorporated herein by reference. Polymerase chain reaction methodologies are well known in the art.

Another method for amplification is the ligase chain reaction ("LCR"), disclosed in EPA No. 320 308, incorporated herein by reference in its entirety. U.S. Pat. No. 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Application No. PCT/US87/00880, incorporated herein by reference, also may be used as still another amplification method in the present invention.

An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[alpha-

thio]-triphosphates in one strand of a restriction site also may be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, i.e., nick translation. A similar method, called Repair Chain Reaction (RCR), involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA. Target specific sequences also can be detected using a cyclic probe reaction (CPR). In CPR, a probe having 3' and 5' sequences of non-specific DNA and a middle sequence of specific RNA is hybridized to DNA that is present in a sample. Upon hybridization, the reaction is treated with RNase H, and the products of the probe identified as distinctive products that are released after digestion. The original template is annealed to another cycling probe and the reaction is repeated.

Still another amplification methods described in GB Application No. 2 202 328, and in PCT Application No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template- and enzyme-dependent synthesis. The primers may be modified by labeling with a capture moiety (e.g., biotin) and/or a detector moiety (e.g., enzyme). In the latter application, an excess of labeled probes are added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS), including nucleic acid sequence based amplification (NASBA) and 3SR Gingeras et al., PCT Application WO 88/10315, incorporated herein by reference.

Davey et al., EPA No. 329 822 (incorporated herein by reference in its entirety) disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention.

Miller et al., PCT Application WO 89/06700 (incorporated herein by reference in its entirety) disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target ssDNA followed by transcription of many RNA copies of the sequence. This scheme is not cyclic, i.e., new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" and "one-sided PCR" (Frohman, 1990 incorporated by reference).

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide", thereby amplifying the di-oligonucleotide, also may be used in the amplification step of the present invention.

Following any amplification, it may be desirable to separate the amplification product from the template and the excess primer for the purpose of determining whether specific amplification has occurred. In one embodiment, amplification products are separated by agarose, agarose-acrylamide or polyacrylamide gel electrophoresis using standard methods. See Sambrook et al., 1989.

Alternatively, chromatographic techniques may be employed to effect separation. There are many kinds of chromatography that may be used in the present invention: adsorption, partition, ion-exchange and molecular sieve, and many specialized techniques for using them including column, paper, thin-layer and gas chromatography.

Amplification products must be visualized in order to confirm amplification of the marker sequences. One typical visualization method involves staining of a gel with ethidium bromide and visualization under UV light. Alternatively, if the amplification products are integrally labeled with radio- or fluorometrically-labeled nucleotides, the amplification products can then be exposed to x-ray film or visualized under the appropriate stimulating spectra, following separation.

In one embodiment, visualization is achieved indirectly. Following separation of amplification products, a labeled nucleic acid probe is brought into contact with the amplified marker sequence. The probe preferably is conjugated to a chromophore but may be radiolabeled. In another embodiment, the probe is conjugated to a binding partner, such as an antibody or biotin, and the other member of the binding pair carries a detectable moiety.

In one embodiment, detection is by Southern blotting and hybridization with a labeled probe. The techniques involved in Southern blotting are well known to those of skill in the art and can be found in many standard books on molecular protocols. See Sambrook et al., 1989. Briefly, amplification products are separated by gel electrophoresis. The gel is then contacted with a membrane, such as nitrocellulose or nylon, permitting transfer of the nucleic acid and non-covalent binding. Subsequently, the membrane is incubated with a chromophore-conjugated probe that is capable of hybridizing with a target amplification product. Detection is by exposure of the membrane to x-ray film or ion-emitting detection devices.

One example of the foregoing is described in U.S. Pat. No. 5,279,721, incorporated by reference herein, which discloses an apparatus and method for the automated electrophoresis and transfer of nucleic acids. The apparatus permits electrophoresis and blotting without external manipulation of the gel and is ideally suited to carrying out methods according to the present invention.

### 5. Expression Constructs

In some embodiments of the present invention, an expression construct that encodes a 3' sequence of GBV-B is utilized. The term "expression construct" is meant to include any type of genetic construct containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a protein, but it need not be. Expression includes both transcription of a gene and translation of mRNA into a gene product. Expression may also include only transcription of the nucleic acid encoding a gene of interest.

In some constructs, the nucleic acid encoding a gene product is under transcriptional control of promoter and/or enhancer. The term promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (tk) and SV40 early transcription units. These studies have shown that promoters are composed of discrete functional modules, each consisting of

approximately 7-20 bp of nucleic acids, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of nucleic acids with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

### 6. Host Cells and Permissive Cells

As used herein, the terms "cell," "cell line," and "cell culture" may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of the present invention, "host cell" refers to a prokaryotic or eukaryotic cell, and it includes any transformable organisms that is capable of replicating a vector or virus and/or expressing viral proteins. A host cell can, and has been, used as a recipient for vectors, including viral vectors. A host cell may be "transfected" or "transformed," which refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny. A "permissive cell" refers to a cell that supports the replication of a given virus and consequently undergoes cell lysis. In the context of the present invention, such a virus would include HCV, GBV-B, or other hepatitis viruses. In a "nonpermissive cell," productive infection does not result, but the cell may become stably transformed. In some embodiments, methods employ permissive cells that are a cell line derived from liver cells (liver cell line).

Host cells may be derived from prokaryotes or eukaryotes, depending upon whether the desired result is replica-

tion of the vector or expression of part or all of the vector-encoded nucleic acid sequences. Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials ([www.atcc.org](http://www.atcc.org)). An appropriate host can be determined by one of skill in the art based on the vector backbone and the desired result. A plasmid or cosmid, for example, can be introduced into a prokaryote host cell for replication of many vectors. Bacterial cells used as host cells for vector replication and/or expression include DH5 $\alpha$ , JM109, and KC8, as well as a number of commercially available bacterial hosts such as SURE $\text{\textcircled{R}}$  Competent Cells and SOLOPACK $\text{\textsuperscript{TM}}$  Gold Cells (STRAT-AGENE $\text{\textcircled{R}}$ , La Jolla). Alternatively, bacterial cells such as *E. coli* LE392 could be used as host cells for phage viruses.

Examples of eukaryotic host cells for replication and/or expression of a vector include HeLa, NIH3T3, Jurkat, 293, Cos, CHO, Saos, and PC12. Many host cells from various cell types and organisms are available and would be known to one of skill in the art. Similarly, a viral vector or virus or virus particle may be used in conjunction with either a eukaryotic or prokaryotic host cell, particularly one that is permissive for replication or expression of the vector. It is contemplated that the present invention includes vectors composed of viral sequences, viruses, and viral particles in the methods of the present invention, and that they may be used interchangeably in these methods, depending on their utility.

Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and their cognate polypeptides, proteins, or peptides.

#### 7. Pharmaceutical Compositions

The present invention encompasses the use of a 3' sequence of GBV-B in the production of or use as a vaccine to combat HCV infection. Compositions of the present invention comprise an effective amount of GBV-B clone as a therapeutic dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. The phrases "pharmaceutically or pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions. For human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

The biological material should be extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle, where appropriate. The active compounds will then generally be formulated for parenteral administration, e.g.,

formulated for injection via the intravenous, intramuscular, sub-cutaneous, intralesional, or even intraperitoneal routes. The preparation of an aqueous composition that contains GBV-B nucleic acid sequences as an active component or ingredient will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and the preparations can also be emulsified.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

A GBV-B clone of the present invention can be formulated into a composition in a neutral or salt form. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. In terms of using peptide therapeutics as active ingredients, the technology of U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, each incorporated herein by reference, may be used.

The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the

required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The preparation of more, or highly, concentrated solutions for direct injection is also contemplated, where the use of DMSO as solvent is envisioned to result in extremely rapid penetration, delivering high concentrations of the active agents to a small area.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above, but drug release capsules and the like also can be employed.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

In addition to the compounds formulated for parenteral administration, such as intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; liposomal formulations; time release capsules; and any other form currently used, including cremes.

Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders. In certain defined embodiments, oral pharmaceutical compositions will comprise an inert diluent or assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 75% of the weight of the unit, or preferably between 25-60%. The amount of active compounds in such therapeutically useful compositions is such that a suitable dosage will be obtained.

The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as mag-

nesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup of elixir may contain the active compounds sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor.

### C. EXAMPLES

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

#### Example 1

##### Elucidation of a 3' Sequence of the GBV-B Genome

The inventors elucidated the previously unrecognized 3' terminal sequence of GBV-B (SEQ ID NO 1). This sequence was reproducibly recovered from tamarin serum containing GBV-B RNA, in RT-PCR nucleic acid amplification procedures using several different primer sets, and as a fusion with previously reported 5' GBV-B sequences.

There is information in the published literature reporting the putative sequences of the 5' and 3' termini of the GBV-B genome. The nucleic acid sequences of these termini were reportedly determined by ligating the ends of the viral RNA together, amplifying the sequence in the region of the resulting junction by reverse-transcription polymerase chain reaction (RT-PCR), and sequencing of the cDNA amplification product across the junction. However, the inventors believed these results required confirmation. Of particular concern was the fact that the 3' terminus appeared to be shorter than the equivalent region of other viruses in the family Flaviviridae (especially within the genus *Hepacivirus*) and that the reported 3' sequence lacked a defined RNA hairpin structure such as those present in these related viruses. Additional novel sequences at the 3' end of the GBV-B genome were investigated using a serum sample collected from a tamarin that was experimentally infected with virus. Amplification was used to determine the sequence of the 3' end.

First, serum (50  $\mu$ L) known to contain GBV-B RNA by RT-PCR assay was extracted with Trizol, and the RNA was washed and dried. A synthetic oligonucleotide was then ligated to the 3' end of the viral RNA. The oligonucleotide, AATTCGGCCCTGCAGGCCACAACAGTC (SEQ ID NO:17), which was phosphorylated at the 5' end and chemically blocked at the 3' end, was ligated to the RNA essentially using the method described by Kolykhalov et al. (Behrens et al., 1996). The RNA was initially dissolved in DMSO and the following additions were made: Tris-Cl, pH

7.5 (10 mM), MgCl<sub>2</sub> (10 mM), DTT (5 mM), hexamine cobalt chloride (1 mM), 10 pmol oligo and 8 U T4 ligase. The final concentration of DMSO was 30% in a final volume of 10 μL. The ligation reaction was incubated for 4 or 20 hours at 19° C. 1 μL of the ligation reaction was used directly to make cDNA, using a primer complementary to the ligated oligonucleotide and the Superscript 2 system, in a final volume of 15 μL. 1 μL of cDNA was amplified using the Advantage cDNA system (Clontech) and two additional oligonucleotide primers. These primers included one that was complementary to the ligated oligonucleotide (i.e., “negative sense”) and a positive-sense primer located near the 3' end of the reported GBV-B sequence. A product approximately 290 bases in length was obtained, and this was gel purified and directly sequenced. Sequencing was done in both directions using the oligonucleotide primers employed for the amplification; 259 bases that had not been previously reported were identified as fused to the sequence that had been previously described as the 3' terminus of the viral genome.

To ensure that this novel 3' sequence from viral RNA could be reproducibly amplified, an additional 10 μL of infected tamarin serum was extracted using Trizol. cDNA was prepared by reverse transcription using an oligonucleotide primer complementary to the penultimate 3' 25 bases of the novel sequence. Amplification was then done by PCR using the primer previously utilized for cDNA synthesis and a positive-sense primer mapping within the previously published GBV-B sequence. In the initial experiments, although a product was readily detected, DNA sequencing showed that this product was missing all of the sequence distal to the poly-U tract. Carrying out the cDNA synthesis in the presence of DMSO circumvented this problem. A cDNA product of approximately 290 bases was obtained. This was sequenced and shown to consist of the 5' primer, 20 bases of the published GBV-B sequence, and 259 bases of the novel sequence obtained in the preceding experiments and containing the sequence of the 3' primer. The sequence of the 3' end of GBV-B is shown in SEQ ID NO: 1 (FIG. 1). The possible secondary RNA structure for this region is shown in FIG. 2, as predicted by a computer-based RNA folding program. The presence of a predicted hairpin structure at the extreme 3' end of this novel sequence is consistent with its location at the 3' terminus of the viral RNA.

The GBV-B cDNA (synthesis described above) was used as a template for PCR amplification of the 3' 1553 nucleotides (nts) of the GBV-B genome. This PCR amplification product was gel purified and cloned into plasmid DNA using the “Perfectly Blunt Cloning Kit” (Novagene).

#### Example 2

##### Construction of an Infectious GBV-B Clone

The elucidation of a 3' sequence of the GBV-B genome will allow those of skill in the art to construct and validate an infectious molecular clone of GBV-B. This will be done using the following procedures.

A full-length cDNA copy of the GBV-B genome containing the newly identified 3' terminal sequences was constructed. RNA transcribed from this cDNA copy of the genome will be infectious when inoculated into the liver of a GBV-B permissive tamarin, giving rise to rescued GBV-B virus particles.

A 1:1000 dilution of GBV-B infectious tamarin serum was obtained. This material was used as a source of viral RNA for the amplification of GBV-B nucleic acid sequences by

reverse-transcription polymerase chain reaction. For amplification of previously reported segments of the GBV-B genome, 250 μL of the diluted serum was extracted with Trizol using the manufacturer's instructions. The final RNA pellet was dissolved in 10 μL of a 100 mM DTT buffer containing 5% RNasin. This material was converted into cDNA using Superscript 2 reverse transcriptase and oligonucleotide primers designed to be complementary to the reported GBV-B RNA sequence and to contain unique restriction sites. This cDNA was amplified using the Advantage cDNA kit (Clontech) employing the cDNA primer (negative sense) as the downstream primer and a similar positive-sense upstream primer, again containing a unique restriction site. The published sequence of GBV-B allowed for the selection of primers in convenient areas of the genome containing unique restriction sites. Using this general strategy, the inventors amplified segments of the reported GBV-B genome representing: (1) nucleotides (nts) 1-1988, using an upstream primer containing a T7 RNA polymerase promoter and a BamHI site upstream of nt 1, and a downstream primer containing a unique EcoRI site (nt 1978); (2) nts 1968-5337, using a downstream primer containing a unique ClaI site at position 5027; (3) nts 5317-7837, using a downstream primer containing a Sall site at nt 7847; and, (4) nts 7837-9143, using a downstream primer containing an added XhoI site. It was found necessary to use different PCR conditions for each primer set.

The RT-PCR products generated in these reactions were cloned into plasmid DNA after gel purification, using the “Perfectly Blunt Cloning Kit” (Novagene). Ten bacterial colonies from each of the four RT-PCR products were analyzed for insert size by restriction endonuclease digestion using EcoRI, the sites for this enzyme being located on either side of the insert in the resulting plasmids. For three of the RT-PCR amplicons, 9 of 10 colonies contained plasmids with the correct size insert. The EcoRI-ClaI amplicon generated only 1/10 colonies with a correct size insert. Thus, 30 additional colonies were examined, yielding two more clones with insert of the correct size. For each of these plasmids, simple restriction patterns were obtained using two restriction enzymes. As these appeared to be correct, the plasmid DNAs were subjected to sequencing using an ABI automatic sequencer.

#### Example 3

##### Nucleotide Sequence of the Cloned GBV-B cDNA

The 5' region of the cloned sequence revealed a relatively long nontranslated region corresponding to the published sequence of the GBV-B 5'NTR, which includes an IRES. This region was followed by a long open reading frame. Near the 3' end of the genome a poly-U tract was identified; however, this was shorter than the published 3' homopolymeric poly-U region. The sequence from these clones was compared with those in the GenBank database (Accession U22304, “Hepatitis GB virus B polypeptide complete genome”). Twenty-two nucleotide differences were identified, of which 14 gave rise to amino acid changes (Table 1). In order to determine whether these changes were genuine or RT-PCR artifacts, which could have been introduced due to the very small amount of material from which these sequences were amplified, segments of the genome containing these changes were reamplified using a serum sample from an independently infected tamarin. Of the 14 changes noted in the original cDNA clones, 12 were not present in these newly amplified sequences and thus were probably

RT-PCR artifacts (Table 2.). A particularly interesting difference from the published GenBank sequence, however, which was present in both the original clones as well as a repeat amplification, was a two-nucleotide substitution that obliterated the SalI site present in the published sequence.

TABLE 1

Differences in the amino acid sequences of GBV-B cDNA clones and the GBV-B sequence reported by Simons et al. (1995).		
GBV-B Protein	AMINO ACID Δ FROM ABBOTT SEQUENCE	RT-PCR Products from Tamarin 12024 (PCR Reaction #)
Core	G <sub>99</sub> →S	G (38.1a)
E1	V <sub>395</sub> →I	V (40.2a)
E2	D <sub>703</sub> →N	D (42.3a)
E2	P <sub>706</sub> →Q	H (42.3a)
E2	A <sub>728</sub> →V	A (42.3a)
NS2	L <sub>791</sub> →F	F (42.3a)
NS2	T <sub>804</sub> →A	A (42.3a)
NS5A	L <sub>1090</sub> →M	L (46.5a)
NS5A	I <sub>2082</sub> →T	I (46.5a)
NS5A	S <sub>2174</sub> →P	(not done)
NS5A	G <sub>2228</sub> →E	E (48.6a)
NS5A	T <sub>2233</sub> →S	T (48.6a)
NS5A	A <sub>2236</sub> →V	A (48.6a)
NS5B	V <sub>2833</sub> →I	V (50.7a)

## Example 4

## Construction of a Full-Length GBV-B cDNA Clone

The four GBV-B cDNA inserts described above were cloned into Bluescript ks+ using unique restriction sites. Since the unique SalI site that was reported to be present in the published GBV-B sequence (nt position 7847) was absent in these cDNA clones, this restriction site was created by engineering two silent nucleotide changes using the "Quick Change" mutagenesis system (Stratagene). Although the most 5' clones (nucleotides 1-7847) could be readily constructed, attempts to add the remaining 3' clones were unsuccessful due to rearrangements and deletions. This problem was overcome by use of pACNR1180, a plasmid that had been used to construct an infectious clone of yellow fever virus. Finally, the most 3' 771 nucleotides of GBV-B were excised from the plasmid containing the novel, previously unreported 3' sequence, and inserted into the truncated assembled GBV-B cDNA construct to complete the 3' end. The 3' terminus of this full-length cDNA was then subjected to DNA sequencing to confirm its integrity. Extensive restriction digests indicated that this construct had the characteristics of a full-length cDNA copy of GBV-B virus. Because there is not yet an understanding of which cultured cells (if any) might be permissive for GBV-B replication, the infectivity of the synthetic GBV-B RNA will be assessed by injecting the RNA directly into the liver of a susceptible tamarin.

Alternatively, an infectious full-length clone can be produced by the following protocol. A plasmid will be made containing a cassette including the 5' and 3' ends of the virus flanked by appropriate restriction sites. These constructs have been shown to efficiently translate reporter genes, with transcription taking place via a T7 promoter placed immediately upstream of the 5'NTR (e.g., see Rijnbrand et al., 1999). The major portion of the GBV-B genome would then be amplified by long range RT-PCR. This method is now well established for hepatitis C virus and other flaviviruses (Teller et al., 1996), and it has been used successfully also to amplify rhinovirus RNA. Briefly this technique uses

"Superscript" reverse transcriptase to synthesize cDNA and a mixture of "KlenTaq 1", and "DeepVent" polymerases to amplify this cDNA. Primers that can be used will contain restriction sites to allow cloning of the RT-PCR products into the cassette vector. After being transformed into suitable competent bacteria, extensive restriction analysis will enable us to determine which clones contain inserts that are of full length and which have a high probability of being correct. Apparent full-length clones will be analyzed further by coupled transcription-translation using the Promega "TnT" system, with the addition of microsomal membranes to allow the cleavage of the structural proteins by cellular signalase enzymes. Clones which appear correct by restriction analysis and which produce GBV-B proteins, in particular the protein coded for by the extreme 3' end of the genome, NS5B, will be selected, and RNA will be transcribed from these clones using the Ambion MegaScript system. "Correct" looking clones (>10) can be injected directly into a tamarin liver at several sites. A successful infection will be determined as described below. If a positive signal is detected the entire genome will be amplified and sequenced to determine which plasmid the virus originated from.

## Example 5

## Rescue of Infectious GVB-B

Infectious GBV-B will be rescued from synthetic genome-length RNA following its injection into the liver of tamarins (*Saquinus* sp.). In past studies, HAV from synthetic RNA in owl monkeys has been recovered (*Aotus trivirgatus*) (Shaffer et al., 1995), and more recently, the recovery of virus from a chimpanzee injected intrahepatically with RNA transcribed from a full-length genotype 1b HCV cDNA clone was reported (Beard et al., 1999).

RNA will be prepared for these experiments using the T7 MegaScript kit (Ambion) and a total of 10 µg of plasmid DNA as template. An aliquot of the reaction products will be utilized to ensure the integrity of the RNA by electrophoresis in agarose-formaldehyde gels. The remainder of the transcription reaction mix will be frozen at -80 C.° until its injection, without further purification, into the liver of a tamarin. Because of the small size of the tamarin, the RNA will be injected under direct visualization following a limited incision and exposure of the liver. Under similar conditions, in other primate species, RT-PCR-detectable viral RNA or cDNA has not been detected in serum samples collected within days of this procedure in the absence of viral replication (Kolykhalov et al., 1997; Yanagi et al., 1998; Beard et al., 1999). Thus, the appearance of RNA in serum collected subsequently from these tamarins will be strong evidence for the replication competence of the synthetic RNA. Serum will be collected weekly for six weeks, then every other week for an additional 6 weeks from inoculated animals. In addition to RT-PCR for detection of viral RNA (see FIG. 2B), alanine aminotransferase (ALT) levels will be measured as an indicator of liver injury and to assess liver histology in punch biopsies taken at the time of ALT elevation. Maximum viremia and an acute phase ALT response is expected to occur around 14-28 days post-inoculation of infectious RNA (Simons et al., 1995; Schlauder et al., 1995; Karayiannis et al., 1989). Transfections will be considered to have failed to give rise to infectious virus if RNA is not detected in the serum within 12 weeks of inoculation. Successfully infected animals will

be followed with twice weekly bleeds until resolution of the viremia, or for 6 months, whichever is longest.

#### Example 6

##### Construction of GBV-B/HCV Chimeras

The GBV-B genome can be used as the acceptor molecule in the construction of chimeric viral RNAs containing sequences of both HCV and GBV-B. Such constructs will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (e.g., proteinase) required for HCV replication. Different classes of chimeric viruses are contemplated. These include: (a) replacement of the GBV-B IRES with that of HCV; and (b) replacement of the NS3 major serine proteinase and helicase, and (c) the replacement of the NS5B RNA-dependent RNA polymerase with the homologous proteins of HCV.

The chimeric constructs described in the following sections will be made by PCR mutagenesis, using high fidelity polymerases and oligonucleotide primers designed to include the specific fusions of GBV-B and HCV sequences (Landt et al., 1990). First round PCR reactions will create the desired fusion, and generate a new "primer" to be used in a second PCR reaction spanning the region to a convenient unique restriction site. PCR cycles will be kept to the minimum number necessary for successful amplification, and all segments of viral sequence that are amplified by PCR will be subjected to DNA sequencing to exclude the presence of unwanted PCR-introduced errors. Sequencing will be accomplished at UTMB's core Recombinant DNA Laboratory. Amplified segments will be kept to the minimum by the exchange of cloned cDNA segments spanning convenient restriction sites in subgenomic clones, and where necessary PCR artifacts can be corrected by site-directed mutagenesis (QuickChange mutagenesis kit, Stratagene).

A number of viable positive-strand RNA virus chimeras have been constructed previously in which IRES elements have been swapped between different viruses. Most of these chimeras have involved the exchange of IRES elements between picornaviruses. Others have been successful in constructing viable poliovirus chimeras containing the HCV IRES in place of the native poliovirus IRES (Zhao et al., 1999; Lu and Wimmer, 1996). A similar rhinovirus 14 chimera containing the HCV IRES has been constructed, although its replication phenotype is not as robust as the poliovirus chimera described by Lu and Wimmer (Lu and Wimmer, 1996). More importantly, Frolov et al. (Frolov et al., 1998) recently reported chimeric flaviviruses in which the HCV IRES was inserted into the genetic background of a pestivirus, bovine viral diarrhea virus (BVDV) in lieu of the homologous BVDV sequence. Although these viable chimeric polioviruses and pestiviruses replicate in cell cultures, they are poor surrogates for HCV in animal models as neither virus is hepatotropic or causes liver disease. Importantly, Frolov et al. (Frolov et al., 1998) demonstrated quite convincingly that the requirement for cis-acting replication signals at the 5' terminus of the pestivirus genome was limited to a short tetranucleotide sequence. This requirement presumably reflects the need for the complement of this sequence at the 3' end of the negative strand during initiation of positive-strand RNA synthesis. The work of Frolov et al.

shows that the IRES of BVDV does not contain necessary replication signals, or that if these are present within the BVDV IRES they can be complemented with similar signals in either the HCV or encephalomyocarditis virus (EMCV, a picornavirus) IRES sequence. Since GBV-B and HCV are more closely related to each other than BVDV and HCV, these studies provide strong support for the viability of chimeras containing the HCV IRES in the background of GBV-B.

Construction of a viable IRES chimera will be enhanced by studies that have documented the sequence requirements and secondary structures of the IRES elements of both HCV and GBV-B (see Lemon and Honda, 1997; Honda et al., 1996; Rijnbrand et al., 1999). To a considerable extent, the work of Frolov et al. (Frolov et al., 1998) was guided by studies of the HCV IRES structure. More recently, these studies have been extended to include a detailed mutational analysis of the GBV-B IRES. The results of these experiments indicate that the functional IRES of GBV-B extends from the 5' end of structural domain II (nt 62) to the initiator AUG codon (nt 446). This segment of the full-length GBV-B clone will be replaced with HCV sequence extending from 5' end of the analogous domain II within the HCV IRES (nt 42) to the initiator codon at the 5' end of the HCV open reading frame (nt 341) to construct the candidate chimera, "GB/C:IRES". The source of HCV cDNA for these studies will be the infectious HCV clone, pCV-H77C, which contains the sequence of the genotype 1a Hutchinson strain virus (Yanagi et al., 1998), whose infectivity in a chimpanzee following intrahepatic inoculation with synthetic RNA transcribed from pCV-H77C has been confirmed.

This GB/C:IRES construct will retain two upstream hairpins within the GBV-B sequence (stem-loops Ia and Ib), and it is thus analogous to the viable "BVDV+HCVdelB2B3H1" chimera of Frolov et al (Frolov et al., 1998). A second chimera can be constructed in which the entire HCV 5' nontranslated RNA will be inserted in lieu of nts 62-446 of the GBV-B genome ("GB/C:5'NTR"). This construct will add to the inserted HCV sequence the most 5' stem-loop from HCV (stem-loop I). A similar insertion was shown to substantially increase the replication capacity of BVDV+HCVdelB2B3H1 by Frolov et al. (Frolov et al., 1998), providing a replication phenotype similar to wild-type BVDV in cell culture.

It is important to point out that there is strong evidence from multiple lines of investigation indicating that it will not be necessary to include coding sequence in these IRES chimeras. This is the case even though Reynolds et al. (Reynolds et al., 1995) have argued that the HCV IRES extends past the initiator codon, and into the core-coding region of that virus. Although Lu and Wimmer (Lu and Wimmer, 1996) found it necessary to include HCV core sequence to obtain a viable chimeric poliovirus, the BVDV chimeras reported by Frolov et al. (Frolov et al., 1998) did not contain any HCV coding sequence. This discrepancy may be explained by the observation that the only downstream requirement for full activity of both the GBV-B and HCV IRES elements is the presence of an unstructured RNA segment (Honda et al., 1996; Rijnbrand et al., 1999). Presumably, this facilitates interaction of the viral RNA with the 40S ribosome subunit in the early steps of cap-independent translation (Honda et al., 1996). The 5' GBV-B coding sequence fulfills this criterion (Rijnbrand et al., 1999).

### In Vitro Characterization of the Translational Activity of IRES Chimeras

The fidelity of the genome-length chimeric constructs will be confirmed by sequencing any DNA segments that have been subjected to PCR during the construction process, as well as confirming sequence at the junction sites. In addition, the translational activity of synthetic RNA derived from these constructs will be assessed and compared to the translational activity of the wild-type GBV-B and HCV RNAs. These studies will be carried out in a cell-free translation assay utilizing rabbit reticulocyte lysates (Rijnbrand et al., 1999). Synthetic RNA will be produced by runoff T7 RNA polymerase transcription using as template ClaI-digested plasmid DNA (BamHI digestion in the case of the genome HCV construct) (T7 Megascript kit, Ambion). <sup>3</sup>H-UTP will be added to the reaction mix to allow for quantification of the RNA product. Reticulocyte lysates (Promega) will be programmed for translation by the addition of RNA (at least 50% full-length as determined by agarose gel electrophoresis) at 20, 40 and 80 µg/ml, and translation reactions will be supplemented with microsomal membranes (Promega). <sup>35</sup>S-Methionine-labelled translation products will be separated by SDS-PAGE, and the quantity of E1 protein produced from each RNA determined by PhosphorImager analysis (Molecular Dynamics). Comparisons of the activity of the HCV IRES in the background of GBV-B and HCV will take into account differences in the methionine content of the E1 proteins of these viruses. Based on previous studies of both the GBV-B and HCV IRES elements (Honda et al., 1996; Rijnbrand et al., 1999), it is expected that these studies will confirm that the HCV IRES will retain nearly full activity when placed within the GBV-B background.

### Example 8

#### In Vivo Characterization of IRES Chimeras

Synthetic RNAs produced from each of the two chimeric GBV-B/HCV constructs (GB/C:IRES and GB/C:5'NTR) will be tested for their ability to induce infection and cause liver disease in susceptible tamarins. These experiments will be carried out as described in Example 2. GB/C:5'NTR may generate viremia and liver injury more closely resembling that observed with wild-type GBV-B infection (Frolov et al., 1998).

### Example 9

#### Chimeric Flaviviruses Containing the HCV NS3 Serine Proteinase/Helicase within the GBV-B Background

Chimeric flaviviruses containing the HCV NS3 serine proteinase/helicase within the GBV-B background are also contemplated within the present invention. The construction of chimeric flaviviruses containing specific heterologous functional polyprotein domains, however, poses a number of special problems. Unlike the situation with the IRES, where the relevant RNA segments appear to have a unique function restricted to cap-independent translation initiation and interact with host cell macromolecules, viral proteins often have multiple functions and may form specific macromolecular complexes with other viral proteins that are essential for

virus replication (Lindenbach and Rice, 1999). Furthermore, such chimeric polyproteins must be amenable to efficient processing by the viral proteinases (NS2/NS3 or NS3). This requires knowledge of the proteinase cleavage specificities as well as specific sites of proteolytic cleavage. Although to date there have been no published studies of the processing of the GBV-B polyprotein, the relatively close relationship between GBV-B and HCV, about 30% overall amino acid identity within the polyprotein (Muerhoff et al., 1995), allows good computer predictions of the alignments of these proteins. The crystallographic structures of both the proteinase and helicase domains of the HCV NS3 protein have been solved (Yao et al., 1997). Thus, both linear alignments and models of the 3D structure of the NS3 proteins of these viruses can provide guidelines for designing specific chimeric fusions that are likely to preserve function.

### Example 10

#### NS3 Proteinase-Domain Chimeras

In HCV, NS3 contains the major serine proteinase that is responsible for most cleavage events in the processing of the nonstructural proteins, i.e., those that occur at the NS3/4A, 4A/4B, 4B/5A and 5A/5B junctions. The active proteinase domain of HCV is located within the amino terminal third of the NS3 protein (residues 1-181), which shares 31% amino acid identity with the analogous segment of the GBV-B polyprotein (GBV-B vs HCV-BK) (Muerhoff et al., 1995). Importantly, the active site of this proteinase appears to be particularly well conserved in GBV-B. The GBV-B proteinase maintains the residues that are responsible for catalysis and zinc binding in the HCV enzyme (Muerhoff et al., 1995), and unlike the NS3 proteinases of some other flaviviruses preserves the Phe-154 residue that determines in part the S<sub>1</sub> specificity pocket of the enzyme and the preference of the HCV proteinase for substrates with a cysteine residue at the P1 position (Scarselli et al., 1997). Thus, it is not surprising that the relevant proteolytic cleavage sites within the GBV-B polyprotein that are predicted from alignments with the HCV polyprotein all possess a Cys residue at this position. Of greatest significance for the proposed experiments, however, is the work of Scarselli et al. (Scarselli et al., 1997) who demonstrated that the GBV-B NS3 proteinase is able to effectively process the polyprotein of HCV in experiments carried out in vitro. Using synthetic peptide substrates, these investigators demonstrated that the enzymatic activities of the GBV-B proteinase (residues 1-181) had kinetic parameters similar to the HCV proteinase on NS4A/4B and NS4B/4A substrates HCV. They did not possess reagents allowing a determination of whether the HCV proteinase is able to cleave a GBV-B substrate, but their results indicate that these viral proteinases share important functional properties. Therefore, these similarities suggest that the HCV proteinase could function in lieu of the GBV-B proteinase if used to replace this segment of an infectious GBV-B clone. In addition, experiments with sindbis/HCV chimeras have shown that the HCV proteinase can cleave within the framework of a sindbis polyprotein (Filocamo et al., 1997). In considering the design of these NS3 proteinase chimeras, there are two additional important considerations. First, in HCV, the cleavage between NS2 and NS3 occurs in cis, as the result of a zinc-dependent metalloproteinase that spans the NS2/NS3 junction (Hijikata et al., 1993). As only the NS3 sequences will initially be exchanged, the viability of the resulting chimeras will be dependent upon preservation of the cis-active cleavage across a chimeric NS2/NS3



proteainase domain. The alignment of GBV-B and HCV sequences shows that residues in HCV that have been shown by Grakoui et al., 1993, to be essential for the NS2/NS3 cleavage are conserved in GBV-B (Muerhoff et al., 1995). Additional chimeras that will include the relevant carboxyl-terminal portion of NS2 can also be created.

A second important consideration is that the mature HCV NS3 proteinase functions as a noncovalent assembly of the NS3 proteinase domain and the amino terminal portion of NS4A, a proteinase accessory factor. The details of this association are well known, and have been studied at the crystallographic level (Kim et al., 1996). The N-terminal domain of the folded proteinase contains eight  $\beta$  strands, including one contributed directly by the NS4A peptide backbone. X-ray studies have shown that this array of  $\beta$  strands gives rise to a much more ordered N-terminus. Thus, the presence of the NS4A strand seems likely to contribute to the structure of the substrate-binding pocket. It is not known whether the NS3 proteinase of GBV-B also requires a similar interaction with NS4A of that virus for complete activity, or, if so, whether the NS4A of GBV-B could substitute for NS4A of HCV in forming the fully active NS3 proteinase of HCV. The predicted GBV-B NS4A molecule is 54 amino acid residues in length (Simons, et al., 1995; Muerhoff et al., 1995), just as in HCV. However, the level of amino acid homology between the NS4A molecules is not especially high, and the potential interaction with either NS3 molecule cannot be predicted from this sequence on the basis of available knowledge. To overcome this potential problem, chimeras will be created in which not only the NS3 proteinase domain of GBV-B is replaced, but also the relevant NS4A segment as well, with homologous segments of the HCV polyprotein. The interaction of the HCV NS3 and NS4A domains represents a unique target for antiviral drug design, and it would be beneficial to have this specific interaction present in any virus to be used as a surrogate for HCV in the evaluation of candidate antiviral inhibitors of HCV proteinase *in vivo*.

The NS3 proteinase chimeras that can be made include "GB/C:NS3<sup>h</sup>", which will contain the sequence encoding the first 181 amino acid residues of the HCV NS3 molecule in lieu of that encoding the first 181 residues of GBV-B NS3, and "GB/C:NS3<sup>h</sup>4A", which will include the same NS3 substitution as well as the HCV sequence encoding the amino-terminal segment of NS4A that forms the interaction with NS3. The precise NS4A sequence to be included in the latter chimera will be based on the modeling studies, which may also suggest more effective fusions of the NS3 proteinase domain of HCV with the downstream NS3 helicase domain of GBV-B. The source of HCV cDNA for these experiments will be the infectious HCV clone, pCV-H77C, which contains the sequence of the genotype 1a Hutchinson strain virus (Yanagi et al., 1998).

#### Example 11

##### NS3 Helicase Domain Chimeras

In addition to serine proteinase activity located within the amino-third of NS3, the downstream carboxy-terminal two-thirds of the molecule contains an RNA helicase activity. These two functional domains appear to be separated by a flexible spacer, within which the fusion of HCV proteinase or helicase domain sequences with GBV-B sequence will be placed. The exact role of the helicase in the HCV life-cycle is not known, but it is almost certainly required for dsRNA strand-separation during some phase of viral RNA synthesis.

The helicase domains of GBV-B and HCV are remarkably well conserved, with some regions within the helicase showing as much as 55% amino acid identity (Muerhoff et al., 1995). The GBV-B helicase is more closely related to the HCV helicase than all other flaviviral NS3 helicases, and it preserves many residues found within the conserved helicase motifs of HCV. Thus the HCV NS3 helicase may be capable of functioning when placed within the polyprotein of GBV-B, and such a chimeric virus may be capable of replication. Residues 182-620 of the GBV-B NS3 molecule will be substituted with the analogous segment of HCV (FIG. 5, "GB/C:NS3<sup>h</sup>"). A chimera will also be made in which the entire NS3 and amino terminal NS4 protein sequences of GBV-B is replaced with the homologous HCV sequences ("GB/C:NS3-4A"). The latter construct will thus represent a dual proteinase-helicase chimera. As with the proteinase chimeras, the HCV cDNA will be derived from pCV-H77C (Yanagi et al., 1998).

#### Example 12

##### In Vitro Characterization of NS3 and NS3-NS4A Chimera

Prior to being evaluated for infectivity in susceptible tamarins, RNAs produced *in vitro* from these clones will be characterized *in vitro*. This evaluation will be restricted to a documentation of the proper processing of the expressed polyprotein (i.e., NS2/NS3 and NS3 proteinase functions), since there are no relevant assays that can determine whether the helicase or RNA-dependent RNA polymerase activities in these polyproteins are sufficient for virus replication. The proteolytic processing of the polyprotein is important, however, as it may be altered either by inclusion of the heterologous HCV NS3 proteinase in lieu of the natural GBV-B protease, or by a change in the folding of the polyprotein induced by inclusion of HCV sequence anywhere within the polyprotein. These studies will be carried out in cell-free coupled transcription/translation assays ("TnT" system, Promega) supplemented with microsomal membranes. Template DNAs will be digested with Sall, which restricts the cDNA within the NS5B coding region. <sup>35</sup>S-methionine-labelled translation products will be separated by SDS-PAGE, and the mature NS3 protein identified by its apparent molecular mass. The NS3 and NS5B proteins will be identified by immunoblot analysis using rabbit antisera to the GBV-B NS3 and NS5B proteins. Generation of a mature ~68 kDa NS3 protein will provide proof of both the cis-active NS2/NS3 cleavage and the NS3-mediated cleavage of NS3/NS4A. Similarly, identification of a mature, processed NS5B molecule will provide further support for the activity of the NS3 proteinase. Controls for these experiments will be the wild-type GBV-B polyprotein expressed in similar fashion from the full-length GBV-B clone. If necessary to more clearly demonstrate the processing of the nonstructural proteins in these constructs, subclones representing the nonstructural region of the chimeric sequences could be produced.

#### Example 13

*In Vivo* Characterization of NS3 and NS3-4A Chimeras  
Synthetic RNAs produced from each of the chimeric GBV-B/HCV constructs described in the preceding section will be tested for their ability to induce infection and cause liver disease in susceptible tamarins. These experiments will be carried out using the approach described above.

#### Chimeric Flaviviruses Containing the HCV NS5B RNA Dependent RNA Replicase within the GBV-B Background

The HCV NS5B molecule contains an RNA-dependent RNA polymerase that plays a central role in replication of the virus. Although this molecule represents a prime target for drug discovery efforts, it has proven difficult to express NS5B in a form that retains enzymatic activity specific for HCV RNA as a substrate. Thus, relatively little is known of the functional activity of the HCV replicase, including structure-function relationships of NS5B. Despite this, the NS5B proteins of GBV-B and HCV appear to be functionally closely related, as they share as much as 43% amino acid identity (Muerhoff et al., 1995). A more important question may be whether an RNA dependent RNA polymerase can act on foreign substrates. However, published work has shown that in vitro purified HCV polymerase has very little specificity for its template, using hepatitis C or globin message with equal fidelity (Behrens et al., 1996; Al et al., 1998). This finding is very similar to that obtained with picornaviral polymerases, where it has been known for many years that in vitro the enzyme exhibits very little specificity. It has always been considered highly likely that this situation would not pertain in vivo where it was thought that the interaction of viral or cellular factors with the 3' end of the genome would generate template specificity. However, recent reports have shown that the removal of the entire 3' untranslated sequence (leaving, however, the poly(A) region present) from both the poliovirus and rhinovirus genome does not completely abrogate the infectivity of the virus (Todd et al., 1997). Furthermore, virus, which was recovered after the initial transfections, was shown to have recovered much of the infectivity of the original virus (Todd et al., 1997). The mechanism for this recovery of infectivity is at present unknown, but these results suggest that the HCV polymerase may be able to function to replicate infectious GBV-B/HCV NS5B chimeras.

Thus a chimeric genome-length virus can be created in which the NS5B coding sequence of HCV (amino acids 2422-3014, 593 residues) is inserted within the background of GBV-B in lieu of its native RNA-dependent RNA polymerase (amino acids 2274-2864, 591 residues). This chimeric virus would be valuable for animal studies of candidate antiviral inhibitors of HCV RNA synthesis.

This NS5B chimera would be evaluated to determine that there was proper proteolytic processing of the polyprotein. This would be accomplished by expression of the chimeric polyprotein in a coupled translation-transcription reaction, followed by immunoblot analysis for the mature NS5B protein, as described for the NS3 and NS3-4A chimeras in the preceding section. If these results confirm that the GB/C:5B chimeric polyprotein is processed with release of NS5B, experiments in tamarins would progress to determine whether synthetic RNA transcribed from the clone is infectious and capable of causing liver disease in intrahepatically inoculated animals. These experiments would be carried out as described above.

A chimeric molecule can be constructed from an infectious GBV-B clone in which the HCV NS3 proteinase or proteinase/helicase sequence would be placed in frame in lieu of the homologous GBV-B sequence, and this chimeric cDNA would be used to generate infectious GBV-B/HCV chimeric viruses by intrahepatic inoculation of synthetic RNA in tamarins. Published studies indicate that the GBV-B and HCV proteinases have closely related substrate recog-

ognition and cleavage properties, likely making such chimeras viable and capable of initiating viral replication in appropriate cell types.

#### Example 15

#### Chimeric Viruses Containing HCV Structural Proteins within a GBV-B Genetic Background, and GBV-B Structural Proteins within an HCV Background

It is well documented that the structural proteins of one flavivirus may in some cases be substituted for those from another member of the family. Such chimeric viruses have been recovered from viruses as distantly related to each other as dengue virus and tick-borne encephalitis virus (Pletnev et al., 1992). More recently, the prM and E proteins of Japanese encephalitis virus have been used to replace the equivalent proteins in a vaccine strain of yellow fever virus to produce a JE/YF chimera (Chambers et al., 1999). These observations suggest that chimeras in which the structural proteins of HCV have been used to replace the homologous proteins of GBV-B may well be viable and capable of replication. The isolation of a chimeric virus containing HCV structural proteins, but having the growth characteristics of GBV-B virus, could answer many fundamental questions concerning the structure and interaction of these proteins in HCV. They would also be useful in addressing the nature of the immune response to HCV structural proteins in infected primates (Farci et al., 1992). More to the point of this application, the availability of such chimeric viruses would allow studies of candidate HCV vaccines to be carried out in the tamarin model. This would be a major advance, because at present such studies are limited to chimpanzees (Choo et al., 1994).

The basis for the difference in the host ranges of HCV and GBV-B is completely unknown. Among many other possibilities, it is conceivable that the host range is dependent upon the availability of a specific receptor(s). If this were the case, host range might be dependent upon the envelope proteins that must interact with the putative cellular receptor. Thus, a chimeric virus containing the envelope proteins of HCV within the genetic background of GBV-B might be noninfectious in tamarins (but potentially infectious in chimpanzees). Thus, a finding that both structural protein chimeras are noninfectious in the tamarin, may require the construction of complementary chimeras in which the relevant GBV-B structural proteins will be inserted into the background of an infectious HCV clone. If inclusion of the GBV-B envelope proteins within the backbone of HCV confers on the resulting chimera the ability to replicate in tamarins, it will confirm an important role for the structural proteins in defining the different host ranges of these viruses. More importantly, the resulting virus would be an exceptionally valuable resource for future studies as it would contain all of the nonstructural replication elements, as well as the 5' and 3' nontranslated regions, of HCV. Such a virus would allow the tamarin model to be used to address many unresolved issues in HCV biology and pathogenesis.

#### Example 16

#### Construction and Evaluation of Structural Protein Chimeras

In designing structural protein chimeras, it is important to note that the two envelope proteins of HCV, E1 and E2, form

## 31

noncovalent heterodimeric complexes that are likely to be important in the assembly of infectious virus particles. This is not known to be the case with the envelope proteins of GBV-B, but it is likely given similarities in the sizes and hydrophathy profiles of these proteins (Simons et al., 1995; Muerhoff et al., 1995). Accordingly, the E1 and E2 proteins will be replaced as a unit, and chimeras containing only one of these proteins from the heterologous virus will generally not be produced. First, a chimera will be created where the E1 and E2 regions of GBV-B virus are replaced with those of HCV, "GB/C:E1-2". The source of HCV cDNA for these constructions will be pCV-H77C (Yanagi et al., 1998). A chimera will also be made in which the core protein, in addition to the envelope proteins, is replaced with the homologous proteins of HCV ("GB/C:Co-E2"). Additional chimeras will be made to determine whether tamarins can be infected with chimeras containing the GBV-B structural proteins within the genetic background of HCV. These will include "C/GB:E1-2" and "C/GB:Co-E2". The backbone for these chimeras will be pCV-H77C, the infectious genotype 1a cDNA clone developed in the Purcell laboratory at NIAID (Yanagi et al., 1998).

The specific amino acid sequences of GBV-B to be replaced with the homologous segments of HCV have been determined by alignments of the GBV-B and HCV sequences, coupled with the location of signalase cleavage sites predicted to be present within the amino terminal third of the GBV-B polyprotein using the computer algorithm of Von Heijne. These predicted signalase cleavages lie between residues 156/157 (core/E1), aa 348-349 (E1/E2) and 732/733 (E2/NS2) in the GBV-B sequence. Thus, the chimera GB/C:E1-2 will contain sequence encoding HCV aa 192-809 in lieu of that encoding aa 157-732 in GBV-B, while the insertion in the GB/C:Co-E2 chimera will extend from the initiator AUG codon (aa 1) to residue 809 in HCV, and will be spliced into GBV-B in lieu of the segment encoding aa 1-732 in the GBV-B clone. The complementary chimeras to be constructed within the background of HCV will involve exchanges of the same segments of the genomes.

## Example 17

## Characterization of Structural Protein Chimeras

Prior to being evaluated for infectivity in tamarins, the processing of these chimeric polyproteins will be examined in coupled transcription/translation reactions supplemented with microsomal membranes, as described in the preceding sections for the proteinase and other proposed chimeras. If these results confirm that the polyprotein is processed as expected, with production of glycosylated E1 and E2 proteins from each of the chimeras (and seen in similar experiments with HCV proteins), studies would proceed in tamarins as previously described. The results of these studies may provide novel information on the basis of the host range differences that exist between HCV and GBV-B. If these results suggest that the envelope proteins play a critical role in determining host range, additional experiments could be carried out with these chimeras in chimpanzees (which are permissive for HCV but apparently not for GBV-B).

## Example 18

## Further Characterization of Rescued Chimeric Viruses

Where infection with chimeric viruses is induced in animals that are injected within the liver with synthetic

## 32

RNA, this virus will be passaged in GBV-B naïve tamarins to further characterize the nature of the infection induced by the chimera. This will be accomplished by taking a pool of the 3 highest titer GBV-B RNA-containing serum specimens from the animal that was successfully transfected with RNA, and inoculating 1 mL of a 1:100 dilution of this pool intravenously into two susceptible animals. These animals will be monitored for infection and liver disease. These animals will be followed until resolution of the viremia and appearance of antibodies detectable in immunoblots with GST-NS3 protein expressed in *E. coli*, or for at least 6 months should an animal sustain a chronic infection. RT-PCR amplification of chimeric segments of the genome may be employed to determine whether the altered phenotype results from mutations within the heterologous portion of the genome.

## Example 19

## Use of GBV-B as Model for HCV

GBV-B and/or GBV-B/HCV chimeras can be used as a model for HCV. Such studies will allow one to investigate the mechanisms for the different biological properties of these viruses and to discover and investigate potential inhibitors of specific HCV activities (e.g., proteinase) required for HCV replication.

## Example 20

## Use of GBV-B/HCV Chimeras to Test Candidate HCV NS3 Proteinase Inhibitors or Other Inhibitors of HCV

GBV-B/HCV viruses can be used in preclinical testing of candidate HCV NS3 proteinase inhibitors or other inhibitors of HCV.

## 1. Candidate Substances

As used herein the term "candidate substance" refers to any molecule that is capable of modulating HCV NS3 proteinase activity or any other activity related to HCV infection. The candidate substance may be a protein or fragment thereof, a small molecule inhibitor, or even a nucleic acid molecule. It may prove to be the case that the most useful pharmacological compounds for identification through application of the screening assay will be compounds that are structurally related to other known modulators of HCV NS3 proteinase activity. The active compounds may include fragments or parts of naturally-occurring compounds or may be only found as active combinations of known compounds that are otherwise inactive. However, prior to testing of such compounds in humans or animal models, it will be necessary to test a variety of candidates to determine which ones have potential.

Accordingly, the active compounds may include fragments or parts of naturally-occurring compounds or may be found as active combinations of known compounds that are otherwise inactive. As such, the present invention provides screening assays to identify agents that are capable of inhibiting proteinase activity in a cell infected with chimeric GBV-B/HCV viruses containing the HCV proteinase. It is proposed that compounds isolated from natural sources, such as animals, bacteria, fungi, plant sources, including leaves and bark, and marine samples may be assayed as candidates for the presence of potentially useful pharmaceutical agents. It will be understood that the pharmaceutical agents to be screened could also be derived or synthesized from chemical compositions or man-made compounds. Thus, it is understood that the candidate substance identified by the present invention may be polypeptide, polynucle-

otide, small molecule inhibitors or any other compounds that may be designed through rational drug design starting from known inhibitors of proteinases or from structural studies of the HCV proteinase.

The candidate screening assays are simple to set up and perform. Thus, in assaying for a candidate substance, after obtaining a chimeric GBV-B/HCV virus with infectious properties, a candidate substance can be incubated with cells infected with the virus, under conditions that would allow measurable changes in infection by the virus to occur. In this fashion, one can measure the ability of the candidate substance to prevent or inhibit viral replication, in relationship to the replication ability of the virus in the absence of the candidate substance. In this fashion, the ability of the candidate inhibitory substance to reduce, abolish, or otherwise diminish viral infection may be determined.

"Effective amounts" in certain circumstances are those amounts effective to reproducibly reduce infection by the virus in comparison to the normal infection level. Compounds that achieve significant appropriate changes in activity will be used. Candidate compounds can be administered by any of a wide variety of routes, such as intravenously, intraperitoneally, intramuscularly, orally, or any other route typically employed.

It will, of course, be understood that all the screening methods of the present invention are useful in themselves notwithstanding the fact that effective candidates may not be found. The invention provides methods of screening for such candidates, not solely methods of finding them.

#### 2. In Vitro Assays

In one particular embodiment, the invention encompasses in vitro screening of candidate substances. Using a cell line that can propagate GBV-B in culture, in vitro screening can be used such that GBV-B or HCV virus production or some indicator of viremia is monitored in the presence of candidate compounds. A comparison between the absence and presence of the candidate can identify compounds with possible preventative and therapeutic value.

#### 3. In Vivo Assays

The present invention also encompasses the use of various animal models to test for the ability of candidate substances to inhibit infection by HCV. This form of testing may be done in tamarins.

The assays previously described could be extended to whole animal studies in which the chimeric virus could be used to infect a GBV-B permissive primate, such as a tamarin. One would then look for suppression of viral replication in the animal, and a possible impact on liver disease related to replication of the infectious chimeric virus. The advantage of this in vivo assay over present available assays utilizing HCV infection in chimpanzees is the reduced cost and greater availability of GBV-B permissive nonhuman primate species.

Treatment of animals with test compounds will involve the administration of the compound, in an appropriate form, to the animal. Administration will be by any route that could be utilized for clinical or non-clinical purposes, including but not limited to oral, nasal, buccal, rectal, vaginal or topical. Alternatively, administration may be by intratracheal instillation, bronchial instillation, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Specifically contemplated are systemic intravenous injection, intraperitoneal injection, and oral administration.

Determining the effectiveness of a compound in vivo may involve a variety of different criteria. Such criteria include, but are not limited to, survival, reduction of rate of infection, arrest or slowing of infection, elimination of infection, increased activity level, improvement in liver function, and improved food intake.

#### Use of Infectious GBV-B/HCV Chimeras as Vaccines

Infectious GBV-B/HCV chimeras expressing HCV envelope proteins will have utility as a vaccine immunogen for hepatitis C. Such clones clearly have the potential to be constructed as chimeras including relevant hepatitis C virus sequences in lieu of the homologous GBV-B sequence, providing unique tools for drug discovery efforts.

Chimeric viruses containing the envelope proteins of hepatitis C virus (as described in the attached) would confer the antigenic characteristics of hepatitis C virus on the chimera. These chimeras may have the ability to replicate in chimpanzees (and thus humans) by virtue of the fact that the chimeric envelope is now able to interact with the human hepatocyte cell surface, a necessary first step in virus replication. Therefore, the chimeric virus, while able to infect and replicate in humans, may not cause much or any disease—the reasoning here is that the genetic backbone of the chimera that encodes the nonstructural proteins of GBV-B has not evolved for replication in human cells and thus may not replicate well. Thus, the chimera may have limited replication ability, cause no disease, but still elicit immunity to the surface envelope proteins of HCV and thus have potential as a hepatitis C vaccine. These chimeras can be tested for their ability to promote immunity to HCV through an immune response.

#### Example 22

#### Construction of a GBV-B Infectious Clone

A genome-length cDNA copy of the complete GB virus B (GBV-B) genome sequence, including the novel 3' terminal sequence, was assembled from fragments amplified by reverse transcription-polymerase chain reaction (RT-PCR) from viral RNA present in a 0.2  $\mu$ l aliquot of infectious serum (GB agent pool mystrax 666, 8/93) supplied by Dr. Jens Bukh of the National Institutes of Health. The serum sample was diluted with 100  $\mu$ l of fetal calf serum and extracted using the Trizol system (GIBCO/BRL). The pellet was dissolved in 10 mM dithiothreitol containing 20 u/mL RNasin (Promega). The selection of primers for cDNA synthesis and PCR amplification was based on the published sequence of GBV-B (Simons et al., 1995). RT/PCR was carried out using Superscript Reverse Transcriptase (GIBCO/BRL) and the Advantage cDNA PCR Kit (Clontech). Four subgenomic regions were amplified covering the entire published GBV-B genome sequence. A fifth subgenomic region included the novel 3' terminal sequence that was identified in our laboratory. The oligonucleotide primer sets (listed 5'→3') used for amplification of the individual regions included:

Primer 1: CGGGATCCCGTAATACGACTCACTATAGACCACAAACACTCCAGTTTG (SEQ ID NO:3)

Primer 2: GTGGAATTCACAGCGTCATA (SEQ ID NO:4)  
(Places a T7 RNA polymerase promoter sequence immediately upstream of the GBV-B sequence; the amplified segment extends from the 5' terminus of the viral genome to the unique EcoRI site at position 1978.)

Primer 3: TGTGAATCCACTCTCTACC (SEQ ID NO:5)

Primer 4: TTATCGATTGCAGCAACCATG (SEQ ID NO:6)

(Overlapping EcoRI site at position 1978 and the unique ClaI site at 5327.)

Primer 5: CATGGTTGCTGCAATCGATAAGCTGAA-GAGTACAATAAC (SEQ ID NO:7)

Primer 6: GACAACAGACGCTTGACACG (SEQ ID NO:8)

(Overlapping the Cla I site at 5327 and a unique Sal I site at 7847 in the published sequence: however, the Sal I site was not present in the amplified GBV-B sequence, and was thus introduced into the cDNA using the QuickChange Site-directed Mutagenesis Kit (Stratagene).)

Primer 7: CTGTCATGGGAGATGCGTAC (SEQ ID NO:9)

Primer 8: CGAGCTCGAGCACATCGCGGGGTCGT-TAAGCCCGGGTCTCC (SEQ ID NO:10)

(Overlapping the Sal I site at 7847 and the published 3' end of the genome.)

Primer 9: GACAACAGACGCTTGACACG (SEQ ID NO:11)

Primer 10: CCGACTCGAGAATTCGGCCCTGCAGGC-CACAACAGTCTCGCGAGTTTTTAATT CCAAGCGGGGTTGCCCTCCGCTTGAA-CAAAAACCGGGGTGCAGCCCTTGG TAC (SEQ ID NO:12)

(Overlaps product of primers 7 and 8 and extends to the novel 3' terminal sequence; includes an Xho I site at the 3' terminus of the GBV-B sequence to aid subsequent manipulations.)

RT-PCR products were gel-purified using the QIAquick Gel Extraction Kit (Qiagen) and ligated into a plasmid vector with the PstBlue-1 Perfectly Blunt Cloning Kit (Novagene). Bacterial colonies were screened for plasmid DNAs with the correct insert size. Those harboring an insert of the approximate correct size were subjected to diagnostic restriction analysis. cDNA inserts generating the appropriate restriction patterns were sequenced in both directions using an ABI DNA sequencer, and those with sequences closest to the published GBV-B sequence (Simons et al., 1995) were selected for subsequent assembly into the full-length clone.

The cloned overlapping cDNA fragments were assembled into the pACNR1180 plasmid using the unique restriction sites noted in the primer descriptions. The final, fully assembled plasmid bearing a 9.4 kb cDNA GBV-B insertion was subjected to DNA sequencing to confirm the validity of the construction and the absence of any mutations or errors in the sequence that may have been introduced during the assembly process. The sequence of the GBV-B infectious cDNA is shown in SEQ ID NO:2.

Plasmid DNA containing the GBV-B cDNA was purified by isopycnic centrifugation on a CsCl gradient and isolated by ethanol precipitation. The plasmid DNA was linearized at the 3' terminus of the cDNA insert by digestion with Xho I. Two µg of the linearized DNA was used as template in an in vitro transcription reaction using the Ambion T7 MEGAscript Kit according to the manufacturer's instructions. The integrity of the resultant synthetic RNA product was confirmed by nondenaturing agarose gel electrophoresis.

Approximately 60 µg of RNA transcribed from the GBV-B cDNA clone was diluted in phosphate buffered saline and injected under direct visualization at laparotomy into the liver of a healthy tamarin without a prior history of GBV-B infection (*Saguinus oedipus*). Serum samples were collected from the animal weekly beginning at 2 weeks post-inoculation and tested for the presence of GBV-B RNA by real-time quantitative RT-PCR (TaqMan assay). Results are shown in the following table.

Week Postinoculation	Genome equivalents/ml
0	0
2	$8.3 \times 10^3$
3	$6.6 \times 10^5$
4	$4.6 \times 10^8$

The presence of high and increasing titers of viral RNA in the serum of this animal between 2-4 weeks postinoculation confirms the infectivity of the full-length cDNA clone containing the novel 3' terminal sequence. The presence of viremia demonstrated that viral replication had ensued following the delivery of the synthetic viral RNA to hepatocytes.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents, which are both chemically and physiologically related, may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Pat. No. 4,578,770  
 U.S. Pat. No. 4,596,792  
 U.S. Pat. No. 4,599,230  
 U.S. Pat. No. 4,599,231  
 U.S. Pat. No. 4,601,903  
 U.S. Pat. No. 4,608,251  
 U.S. Pat. No. 4,683,195  
 U.S. Pat. No. 4,683,202  
 U.S. Pat. No. 4,800,159  
 U.S. Pat. No. 4,883,750  
 U.S. Pat. No. 5,279,721  
 U.S. Pat. No. 5,807,670  
 EPA No. 320,308  
 EPA No. 329,822  
 GB Application No. 2,202,328  
 PCT/US87/00880  
 PCT/US89/01025  
 WO 88/10315  
 WO 89/06700  
 WO 90/07641  
 Al, R H., Xie, Y., Wang, Y., and Hagedon: Expression of recombinant hepatitis C NS5B in *E. coli*. *Virus Res* 53 141-149, 1998.  
 Alter, M. J. Epidemiology of hepatitis C. *Hepatology* 26 625-655, 1997.  
 Beard, M R., Abell, G., Sangar, D V., Honda, M., Garland, M., Carroll, A R., and Lemon S M: An Infectious Molecular Clone of a Japanese Genotype 1b Hepatitis C Virus. *Hepatology, In press*, 1999.

Behrens, S E., Tomei, L., and De Francesco: Identification and properties of the RNA dependent RNA polymerase of hepatitis C virus. *EMBO J* 15 12-22, 1996.

Chambers, T J, Nestorowicz, A., Mason, P W., and Rice, C M. Yellow fever/Japanese encephalitis chimeric viruses. Construction and biological properties. *J Virol* 73 3095-3101, 1999.

Choo, Q.-L., G. Kuo, G. Ralston, R., Weiner, A., Chien, D., VanNest, G., Han, P., Berger, K. Thudium, K., Kuo, C., Kansopon, J., McFarland, J., Tabrizi, A., Ching, K., Moss, B. Cummins, L. B., Houghton, M. and Muchmore, E., Vaccination of chimpanzees against infection by the hepatitis C virus. *Proc Natl Acad Sci USA* 91:1294-1298, 1994.

Consensus Development Panel. National Institutes of Health Consensus Development Conference Panel statement: Management of hepatitis C. *Hepatology* 26, Supplement 1[3], 2S-10S, 1997.

Farci, P., Alter, H. J., Govindarajan, S., Wong, D. C., Engle, R., Lesniewski, R. R., Mushahwar, I. K., Desai, S. M., Miller, R. H., Ogata, N., and Purcell, R. H., Lack of protective immunity against reinfection with hepatitis C virus. *Science* 258 135-140, 1992.

Farci, P., London, W. T., Wong, D. C., Dawson, G. J., Vallari, D. S., Engle, R., and Purcell, R. H., The natural history of infection with hepatitis C virus (HCV) in chimpanzees: Comparison of serologic responses measured with first- and second-generation assays and relationship to HCV viremia. *The Journal of Infectious Diseases* 165, 1006-1011, 1992.

Filocamo, G., Pacinsh, L., and Migliacciog: Chimeric sindbis virus dependent on the NS3 protease of hepatitis C virus. *J Virol* 71 1417-27, 1997.

Frohman, *PCR PROTOCOLS: A GUIDE TO METHODS AND APPLICATIONS*, Academic Press, New York, 1990.

Frolov, I., McBride, M. S., and Rice, C. M., cis-acting RNA elements required for replication of bovine viral diarrhea virus-hepatitis C virus 5' nontranslated region chimeras. *RNA* 4, 1418-1435, 1998.

Grakoui et al., Characterization of the hepatitis C virus-encoded serine proteinase: determination of proteinase-dependent polypeptide cleavage sites. *J. Virol.* 67(5): 2832-43, 1993.

Grakoui et al., Expression and identification of hepatitis C virus polyprotein cleavage products. *J. Virol* 67(3):1385-95, 1993.

Hijikata, M., Mizushima, H., Akagi, T., Mori, S., Kakiuchi, N., Kato, N., Tanaka, T., Kimura, K., and Shimotohno, K: Two distinct protease activities required for the processing of a putative non-structural precursor protein of hepatitis C virus. *J Virol* 67 4665-4675, 1993.

Honda, M., Brown, E A., and Lemon, S M: Stability of a stem-loop involving the initiator AUG controls the efficiency of internal initiation of translation on hepatitis C virus RNA. *RNA* 2 955-956, 1996.

Karayiannis, P., Petrovic, L M., Fry, M., Moore, D., Enticott, M., McGarvey, M J., Scheuer, P J., and Thomas, H C: Studies of GB hepatitis agent in tamarins. *Hepatology* 9 186-192, 1989.

Kim, J. L., Morgenstern, K. A., Lin, C., Fox, T., Dwyer, M. D., Landro, J. A., Chambers, S. P., Markland, W., Lepre, C. A., O'Malley, E. T., Harbeson, S. L., Rice, C. M., Murcko, M. A., Caron, P. R., and Thomson, J. A.: Crystal structure of the hepatitis C virus NS3 protease domain complexed with a synthetic NS4A cofactor peptide. *Cell* 87, 343-355, 1996.

Kolykhalov, A A., Agapov, E N., Blight, K., Mihalik, S M., Feinstone, S M., and Rice, C M: Transmission of hepatitis C by intrahepatic inoculation of transcribed RNA. *Science* 277 570-574, 1997.

5 Kolykhalov, A A., Feinstone, S M., and Rice, C M: Identification of a highly conserved element at the 3' terminus of hepatitis C virus genome RNA. *J Virol* 70 3363-3371, 1996.

Landt, O., Grunert, H. P., Hahn, U. A general method for rapid site-directed mutagenesis using the polymerase chain reaction. *Gene* 96: 125-128, 1990.

Lemon, S. M. and Honda, M. (1997) Internal ribosome entry sites within the RNA genomes of hepatitis C virus and other flaviviruses. *Seminars in Virology* 8, 274-288, 1997.

15 Lindenbach, B. D. and Rice, C. M. Genetic interaction of flavivirus nonstructural proteins NS1 and NS4A as a determinant of replicase function [In Process Citation]. *J Virol* 73, 4611-4621, 1999.

Lu, H.-H. and Wimmer, E. Poliovirus chimeras replicating under the translational control of genetic elements of hepatitis C virus reveal unusual properties of the internal ribosomal entry site of hepatitis C virus. *Proc Natl Acad Sci USA* 93, 1412-1417, 1996.

Muerhoff, A S., Leary, T P., Simons, J N., Pilot-Matias, T J., Dawson, G J., Erker, J C., Chalmers, M L., Schlauder, G G., Desai, S M., and Mushahwar, I K: Genomic organization of GB viruses A and B: Two new members of the flaviviridae associated with GB agent hepatitis. *J Virol* 69 5621-5630, 1995.

25 Neumann, A. U., Lam, N. P., Dahari, H., Gretch, D. R., Wiley, T. E., Layden, T. J., and Perelson, A. S. Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science* 282, 103-107, 1998.

Pletnev, A. G., Bray, M., Huggins, J. and Lai, C. J. Construction and characterization of chimeric tick-borne encephalitis and dengue viruses. *Proc. Natl. Acad Sci USA* 89; 10532-10536, 1992.

Remington's Pharmaceutical Sciences, 15<sup>th</sup> Edition, pages 1035-1038 and 1570-1580.

40 Reynolds, J. E., Kaminiski, A., Kettinen, H. J., Carroll, A. R., Rowlands, D. J., and Jackson, R. J. Unique features of internal initiation of hepatitis C virus RNA translation. *EMBO J.* 14, 6010-6020, 1995.

Rijnbrand, R., Abell G, and Lemon, S. M. Mutational analysis of the GB virus B internal ribosome entry site. Submitted for publication, 1999.

Sambrook, et al., *Molecular Cloning, A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1989.

50 Scarselli, E., Urbani, A., Sbardellati, A., Tomei, L., De Francesco, R., and Traboni: GB virus B and hepatitis C virus share substrate specificity. *J Virol* 71 4985-4989, 1997.

Schlauder, G G., Dawson, G J., Simons, J M., Pilot-Matias, T J., Gatierez, R A., Heynen, C A., Knigge, M F., Kurpiewski, G S., Buijk, S L., Leary, T P., Muerhoff, A S., Desai, S M., and Mushahwar, I K; Molecular and serological analysis in the transmission of the GB hepatitis agent. *J Med Virol* 46 81-90, 1995.

60 Shaffer, D. R., Emerson, S. U., Murphy, P. C., Govindarajan, S., and Lemon, S. M. A hepatitis A virus deletion mutant which lacks the first pyrimidine-rich tract of the 5' non-translated RNA remains virulent in primates after direct intrahepatic nucleic acid transfection. *J Virol* 69, 6600-6604, 1995.

Simons, J N., Pilot-Matias, T J., Leary, S L., Dawson, G J., Desai, S M., Schlauder, G G., Muerhoff, A S., Erker, J C.,

- Buijk, S L., Chalmers, M L., Van Saint, C L., and Mushahwar, I K: Identification of two flavivirus like genomes in the GB hepatitis agent. *Proc Natl Acad Sci* 92 3401-3405, 1995.
- Tanaka, T., Kato, N., Cho, M.J., and Shimothono, K: A novel sequence found at the 3' terminus of hepatitis C virus genome. *Biochem Biophys Res Comm* 215 744-749, 1995.
- Teller, R., Bukh, J., Emerson, S.U., Miller, R H., and Purcell, R H., Long PCR and it's application to hepatitis A, hepatitis B and hepatitis C virus genomes. *J Clin Micro* 34 3085-3091, 1996.
- Todd, S., Towner, J S., Brown, DM., and Semler, B L: Replication competent picornaviruses with complete genomic RNA 3' non-coding deletions. *J Virol* 71 8868-74, 1997.

- Yanagi, M., Purcell, R H., Emmerson, S U., and Bukh, J., Transcripts from a single full length clone are infectious when directly transfected into a liver of a chimpanzee. *Proc Natl Acad Sci USA* 16 8738-8743, 1998.
- 5 Yao, N. H., Hesson, T., Cable, M., Hong, Z., Kwong, A. D., Le, H. V., and Weber, P. C., Structure of the hepatitis C virus RNA helicase domain. *Nature Structural Biology* 4, 463-467, 1997.
- 10 Zhao, W D., Wimmer, E., and Lahser, F C. Poliovirus/hepatitis C virus (internal ribosome entry site-core) chimeric viruses: improved growth properties through modification of a proteolytic cleavage site and requirement for core RNA sequence but not for core related polypeptides. *J Virol* 73 1546-1554, 1999.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1  
 <211> LENGTH: 260  
 <212> TYPE: RNA  
 <213> ORGANISM: Unknown  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Unknown Organism: GB VIRUS B

<400> SEQUENCE: 1

```
gaguuuugcga ccauggugga ucagaaccgu uucgggugaa gccauggucu gaaggggaug      60
acgucccuuc uggucaucc acaaaaaccg ucucgggugg gugaggaguc cuggcugugu      120
gggaagcagu caguuaauu cccgucgugu guggugacgc cucacgacgu auuuguccgc      180
ugugcagagc guaguaccaa gggcugcacc ccgguuuuug uuccaagcgg agggcaaccc      240
ccgcuuggaa uaaaaaacug                                     260
```

<210> SEQ ID NO 2  
 <211> LENGTH: 9399  
 <212> TYPE: DNA  
 <213> ORGANISM: GBV-A-like virus

<400> SEQUENCE: 2

```
accacaaaca ctccagtttg ttacactcgg ctaggaatgc tcttgagca cccccctag      60
cagggcgtgg gggatttccc ctgccctctc gcagaagggt ggagccaacc accttagtat      120
gtaggcggcg ggactcatga cgctcgcgtg atgacaagcg ccaagcttga cttgatggc      180
cctgatgggc gttcatgggt tcggtggtgg tggcgcttta ggcagcctcc acgcccacca      240
cctcccagat agagcggcgg cactgtaggg aagaccgggg accggtcact accaaggacg      300
cagacctctt tttgagtatc acgcctccgg aagtagttgg gcaagccac ctatatgtgt      360
tgggatggtt ggggttagcc atccataccg tactgcctga tagggtcctt gcgaggggat      420
ctgggagtct cgtagaccgt agcacatgcc tgttatttct actcaacaa gtcctgtacc      480
tgcgcccaga acgcgcaaga acaagcagac gcaggcttca taccctgtgt cattaataaac      540
atctgttgaa aggggacaac gagcaaagcg caaagtccag cgcgatgctc ggcctcgtaa      600
ttacaaaatt gctggtatcc atgatggcct gcagacattg gctcaggctg ctttgccagc      660
tcatggttgg ggacgccaag accctcgcca taagtctcgc aatcttgtaa tcttctgga      720
ttaccctttg gggtgattg gtgatgttac aactcacaca cctctagtag gcccgctgg      780
ggcaggagcg gtcgctcgac cagtctgcca gatagtacgc ttgctggagg atggagtcaa      840
```

-continued

---

ctgggctact	ggttggttcg	gtgtccacct	ttttgtgta	tgtctgctat	ctttggcctg	900
tcccgttagt	ggggcgggg	tcactgaccc	agacacaaat	accacaatcc	tgaccaattg	960
ctgccagcgt	aatcaggtta	tctattgttc	tccttccact	tgcttacacg	agcctggttg	1020
tgtgatctgt	gcgagcaggt	gctgggttcc	cgccaatccg	tacatctcac	acccttccaa	1080
ttggactggc	acggactcct	tcttggttga	ccacattgat	tttgttatgg	gcgctcttgt	1140
gacctgtgac	goccttgaca	ttggtgagtt	gtgtggtgcg	tgtgtattag	tcggtgactg	1200
gcttgcagg	cactggctta	ttcacataga	cctcaatgaa	actggtactt	gttacctgga	1260
agtgcccact	ggaatagatc	ctgggttccct	agggtttata	gggtggatgg	ccggcaaggt	1320
cgaggctgtc	atcttcttga	ccaaactggc	ttcacaagta	ccatacgeta	ttgogactat	1380
gttttagcagt	gtacactacc	tggcggttgg	cgctctgata	tactatgcct	ctcggggcaa	1440
gtggtatcag	ttgctcctag	cgcttatgct	ttacatagaa	gcgacctctg	gaaaccctat	1500
cagggtgccc	actggatgct	caatagctga	gttttgctcg	cctttgatga	taccatgtcc	1560
ttgccactct	tatttgagtg	agaatgtgtc	agaagtatt	tgttacagtc	caaagtggac	1620
caggcctatc	actctagagt	ataacaactc	catacttgg	taccctata	caatccctgg	1680
tgcgagggga	tgtatggtta	aattcaaaaa	taacacatgg	ggttgctgcc	gtattcgcaa	1740
tgtgccatcg	tactgacta	tgggactga	tgacgtgtgg	aacgacactc	gcaaacctta	1800
cgaaagtatg	ggtgtaaac	catggctaac	aaccgcatgg	cacaacggct	cagccctgaa	1860
attggtata	ttacaatacc	ctgggtctaa	agaaatgttt	aaacctata	attggatgtc	1920
aggccatttg	tattttgagg	gatcagatac	ccctatagtt	tacttttatg	accctgtgaa	1980
ttccactctc	ctaccaccgg	agaggtgggc	taggttgccc	ggtacccac	ctgtggtacg	2040
tggttcttgg	ttacaggttc	cgcaaggttt	ttacagtgat	gtgaaagacc	tagccacagg	2100
attgatcacc	aaagcaaaa	cctggaaaaa	ttatcaggtc	ttatattccg	ccacgggtgc	2160
tttctctctt	acgggagtta	ccaccaaggc	cggtgtgcta	attctgttgg	ggttgtgtgg	2220
cagcaagtat	cttattttag	cctacctctg	ttacttgtcc	ctttgttttg	ggcgcgcttc	2280
tggttaccct	ttgcctcctg	tgctcccatc	ccagtcgtat	ctccaagctg	gctgggatgt	2340
tttctctaaa	gctcaagtag	ctccttttgc	tttgattttc	ttcatctggt	gctatctccg	2400
ctgcaggcta	cgttatgctg	cccttttagg	gtttgtgccc	atggctgceg	gcttgcccct	2460
aactttcttt	ggtgcagcag	ctgctgccc	accagattat	gactggtggg	tgcgactgct	2520
agtggcaggg	ttagttttgt	gggcccggcg	taaccgtggt	caccgcatag	ctctgcttgt	2580
aggtccttgg	cctctggtag	cgcttttaac	cctcttgcat	ttggttacgc	ctgcttcagc	2640
ttttgatacc	gagataaatt	gagggtgac	aataccacct	gtagtagcat	tagttgtcat	2700
gtctcgtttt	ggcttcttgg	ctcacttgtt	acctcgctgt	gctttagtta	actcctatct	2760
ttggcaacgt	tgggagaatt	ggttttgaa	cgttacacta	agaccggaga	ggtttttcct	2820
tgtgctgggt	tgtttcccgg	gtgcgacata	tgacgcgctg	gtgactttct	gtgtgtgca	2880
cgtagctctt	ctatgtttaa	catccagtc	agcatcgttc	tttgggactg	actctaggg	2940
tagggcccat	agaatgttgg	tgctctcgg	aaagtgtcat	gcttggatt	ctcattatgt	3000
tcttaagttt	ttcctcttag	tgtttgggtga	gaatgggtgtg	tttttctata	agcacttgca	3060
tggtgatgtc	ttgcctaatt	atthtgcctc	gaaactacca	ttgcaagagc	catttttccc	3120
ttttgaaggc	aaggcaaggg	tctatagaa	tgaaggaaga	cgcttggcgt	gtggggacac	3180



-continued

---

ggttgatggt ttgcccggtg ttgcgcgtct cggcgacctt gttttcgag ggttgctat	3240
gccgccagat ggggtggcca ttaccgcacc ttttacgctg cagtgtctct ctgaacgtgg	3300
cacgctgtca gcgatggcag tggcatgac tggtatagac cccggaactt ggactggaac	3360
tatcttcaga ttaggatctc tggccactag ctacatggga tttgtttgtg acaacgtgtt	3420
gtatactgct caccatggca gcaagggcg cgggttgct catcccacag gctctataca	3480
cccaataaacc gttgacgcgg ctaatgacca ggacatctat caaccacat gtggagctgg	3540
gtcccttact cgggtctctt gcggggagac caaggggtat ctgtaaacac gactggggtc	3600
attggttgag gtcaacaaat ccgatgacct ttattggtgt gtgtgcgggg cccttcccat	3660
ggctgttgcc aagggttctt cagggtcccc gattctgtgc tcctccgggc atgttattgg	3720
gatgttcacc gctgctagaa attctggcgg ttcagtcagt cagattaggg ttaggccgtt	3780
ggtgtgtgct ggataccatc cccagtacac agcacatgcc actcttgata caaacctac	3840
tgtgcctaac gagtattcag tgcaaat tttt aattgcccc actggcagcg gcaagtcaac	3900
caaattacca ctttcttaca tgcaggagaa gtatgaggtc ttggtcctaa atcccaggt	3960
ggctacaaca gcatcaatgc caaagtacat gcacgcgacg tacggcgtga atccaaattg	4020
ctattttaat ggcaaatgta ccaacacagg ggcttcactt acgtacagca catatggcat	4080
gtacctgacc ggagcatgtt cccggaacta tgatgtaatc atttgtgacg aatgccatgc	4140
taccgatgca accaccgtgt tgggcattgg aaaggtccta accgaagtc catccaaaa	4200
tgtaggcta gtggttcttg ccacggctac cccccctgga gtaatcccta caccacatgc	4260
caacataact gagattcaat taaccgatga aggcactatc ccctttcatg gaaaaaagat	4320
taaggaggaa aatctgaaga aaggagaca ccttatcttt gaggtacca aaaaactg	4380
tgatgagctt gtaacagagt tagctcgaaa ggaataaca gctgtctctt actataggg	4440
atgtgacatc tcaaaaatcc ctgagggcga ctgtgtagta gttgcaactg atgccttg	4500
tacagggtag actggtgact ttgattccgt gtatgactgc agcctcatgg tagaaggcac	4560
atgccatgtt gacctgacc ctactttcac catgggtgtt cgtgtgtgcg gggtttcagc	4620
aatagttaaa ggccagcgtg gggcgcgac aggccgtggg agagctggca tatactacta	4680
tgtagacggg agttgtacc cttcgggtat ggttctgaa tgcaacattg ttgaagcctt	4740
cgacgcagcc aaggcatggt atggtttgc atcaacagaa gctcaacta ttctggacac	4800
ctatgcacc caacctgggt tacctgcgat aggagcaaat ttggacgagt gggctgatct	4860
cttttctatg gtcaacccc aacctcatt tgtcaatact gcaaaaagaa ctgctgacaa	4920
ttatgtttt ttgactgcag cccaactaca actgtgtcat cagtattggt atgctgctcc	4980
caatgacgca ccacggtggc agggagccc gcttgggaaa aaacctgtg gggttctgtg	5040
gcgcttgac gccgctgac cctgtcctgg ccagagccc agcgagggtg ccagatacca	5100
aatgtgcttc actgaagtca atacttctgg gacagccgca ctgctgttg gcgttgag	5160
ggctatggct tatctagcca ttgacactt tggcgccact tgtgtgcggc gttgctggtc	5220
tattgatca gtccctacc gtgctactgt cgcgccagtg gttgacgaag aagaaatcgt	5280
ggaggagtgt gcatcattca ttcccttggg ggccatggt gctgcaatcg ataagctgaa	5340
gagtacaatc accacaacta gtcctttcac attggaaacc gccctgaaa aacttaacac	5400
ctttcttgg cctcatgcag ctacaatcct tgctatcata gagtattgct gtggtttagt	5460
cactttacct gacaatccct ttgcatcatg cgtgtttgct ttcattgcgg gtattactac	5520
cccactacct cacaagatca aatgttctc gtcattat ttt ggaggcga ttgogtccaa	5580

-continued

---

gcttacagac gctagaggcg cactggcggtt catgatggcc ggggctgcgg gaacagctct	5640
tggtacatgg acatcgggtg gttttgtctt tgacatgcta ggcggctatg ctgccgctc	5700
atccaactgct tgcttgacat ttaaagtctt gatgggtgag tggcccacta tggatcagct	5760
tgctggttta gtctactccg cgttcaatcc ggccgcagga gttgtgggcg tcttgcagc	5820
ttgtgcaatg tttgctttga caacagcagg gccagatcac tggccaaca gacttcttac	5880
tatgcttgct aggagcaaca ctgatatgaa tgagtacttt attgccaactc gtgacatccg	5940
caggaagata ctgggcatc tggaggcatc taccctctgg agtgcatac cagcttgcat	6000
ccgttggtc cacaccccg cggaggatga ttgcggctc attgcttggg gtctagagat	6060
ttggcagtat gtgtgcaatt tctttgtgat ttgctttaat gtccttaaag ctggagttca	6120
gagcatgggt aacattcctg gttgtccttt ctacagctgc cagaaggggt acaagggccc	6180
ctggattgga tcaggatgac tccaagcagc ctgtccatgc ggtgctgaac tcatcttttc	6240
tggtgagaat ggttttgcaa aactttacaa aggaccaga acttgttcaa attactggag	6300
aggggtggt ccagtcacg ctaggctgtg tgggtcggct agaccggacc caactgattg	6360
gactagtctt gtcgcaatt atggcgtag ggactactgt aaatatgaga aaatgggaga	6420
tcacatttt gttacagcag tatcctctcc aaatgtctgt ttcaccagc tgcaccaac	6480
cttgagagct gcagtgccg tggacggcgt acaggttcag tgttatctag gtgagcccaa	6540
aactccttg acgacatctg ctgtctgta cggctctgac ggtaaaggta aaactgttaa	6600
gcttccctc cgcgttgac gtccacacacc tgggtgctgc atgcaacta atttgcgtga	6660
tgcaactgag acaaatgact gtaattccac aaacaacact cctagtgatg aagccgcagt	6720
gtccgctctt gttttcaaac aggagttgcg cgttacaac caatgcttg aggcaatttc	6780
agctggcgtt gacaccacca aactgccagc cccctccatc gaagagtag tggtaagaaa	6840
gcgccagttc cgggcaagaa ctggttcgct taccttgct cccctccga gatccgtccc	6900
aggagtgtca tgtcctgaaa gcctgcaacg aagtgaccg ttagaaggtc cttcaaacct	6960
ccctccttca ccactgttc tacagttgac catgccgatg cccctgttg gagcgggtga	7020
gtgtaaccct tcaactgcaa ttggatgtc aatgaccgaa acagcggag gccctgatga	7080
tttaccagc taccctccca aaaaggaggc ctctgaatgg tcagacgaaa gttggtcagc	7140
ggctacaacc gtttccagct acgttactgg cccccgtac cctaagatac ggggaaagga	7200
ttccactcag tcagccccg ccaaacggcc tacaaaaag aagttgggaa agagtgagtt	7260
ttcgtgcagc atgagctaca cctggaccga cgtgattagc tcaaaaactg cttctaaagt	7320
tctgtctgca actcgggcca tcaactagtg tttcctcaa caaagatcat tgggtatgt	7380
gactgagccg cgggatgcg agcttagaaa acaaaaagtc actattaata gacaacctct	7440
gttccccca tcataccaca agcaagtgag attggctaag gaaaaagctt caaaagttgt	7500
cgggtgcatg tgggactatg atgaagtagc agctcacacg ccctctaagt ctgctaagtc	7560
ccacatcaact gcccttcggg gcaactgatg tcgttctgga gcagcccga aggctgttct	7620
ggacttgagc aagtgtgtgc aggcaggtga gataccgagc cattatcggc aaactgtgat	7680
agttccaaag gaggaggtct tcgtgaagac ccccagaaa ccaacaaaga acccccaag	7740
gcttatctcg taccctccac ttgaaatgag atgtgttgag aagatgtact acggtcaggt	7800
tgctcctgac gtagttaaag ctgtcatggg agatgcgtac gggttttag atccacgtac	7860
ccgtgtcaag cgtctgttgt cgatgtggtc acccgatgca gtcggagcca catgcgatac	7920

-continued

---

```

agtggtgtttt gacagtacca tcacacccga ggatatcatg gtggagacag acatctactc 7980
agcagctaaa ctcaatgacc aacaccgagc tggcattcac accattgcga ggcagttata 8040
cgctggagga ccgatgatcg cttatgatgg ccgagagatc ggatatcgta ggtgtaggtc 8100
ttccggcgtc tatactacct caagttccaa cagtttgacc tgctggctga aggtaaagtc 8160
tgacgccgaa caggctggca tgaagaacct tcgcttcctt atttgcggcg atgattgcac 8220
cgtaatttgg aagagcgccg gagcagatgc agacaaacaa gcaatgcgtg tctttgctag 8280
ctggatgaag gtgatgggtg caccacaaga ttgtgtgcct caacccaaat acagtttgga 8340
agaattaaca tcatgctcat caaatgttac ctctggaatt accaaaagtg gcaagcctta 8400
ctactttcctt acaagagatc ctctatccc ccttggcagg tgctctgccg agggctctgg 8460
atacaacccc agtgcgtcgt ggattgggta tctaatacat cactaccat gtttgggg 8520
tagccgtgtg ttggctgtcc atttcatgga gcagatgctc tttgaggaca aacttcccga 8580
gacggtgacc tttgactggt atgggaaaaa ttatacggtg cctgtagaag atctgccag 8640
catcattgct ggtgtgcacg gtattgaggc tttctcggtg gtgcgtaca ccaacgctga 8700
gatcctcaga gtttcccaat cactaacaga catgaccatg cccccctgc gagcctggcg 8760
aaagaaagcc agggcggtcc tcgccagcgc caagaggcgt ggcggagcac acgcaaaatt 8820
ggctcgttc cttctctggc atgctacatc tagacctcta ccagatttg ataagacgag 8880
cgtggctcgg tacaccactt tcaattattg tgatgtttac tccccggagg gggatgtgtt 8940
tattacacca cagagaagat tgcagaagtt tcttgtgaag tatttggctg tcattgtttt 9000
tgccctaggg ctcatgctg ttggattagc catcagctga acccccaaat tcaaaattaa 9060
ctaacagttt tttttttttt tttttttttt agggcagcgg caacagggga gaccccgggc 9120
ttaacgacct cgccgatgtg agtttggcga ccatgggtgga tcagaaccgt ttcgggtgaa 9180
gccatggtct gaaggggatg acgtcccttc tggtcatcc acaaaaaccg tctcgggtgg 9240
gtgaggagtc ctggctgtgt gggaagcagt cagtataatt cccgctcgtg gtggtgacgc 9300
ctcacgacgt atttgcgcg tgtgcagagc gtagtaccaa gggctgcacc ccggttttt 9360
ttccaagcgg agggcaaccc ccgcttgaa ttaaaaact 9399

```

```

<210> SEQ ID NO 3
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

```

```
<400> SEQUENCE: 3
```

```
cgggatcccg taatacgact cactatagac cacaacact ccagtttg 48
```

```

<210> SEQ ID NO 4
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

```

```
<400> SEQUENCE: 4
```

```
gtggaattca cagcgtcata 20
```

```

<210> SEQ ID NO 5
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 5  
 tgtgaattcc actctcctac c 21

<210> SEQ ID NO 6  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 6  
 ttatcgattg cagcaaccat g 21

<210> SEQ ID NO 7  
 <211> LENGTH: 39  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 7  
 catggttgct gcaatcgata agctgaagag tacaataac 39

<210> SEQ ID NO 8  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 8  
 gacaacagac gcttgacacg 20

<210> SEQ ID NO 9  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 9  
 ctgtcatggg agatcgctac 20

<210> SEQ ID NO 10  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 10  
 cgagctcgag cacatcgcg ggctgtaag cccgggtct cc 42

<210> SEQ ID NO 11  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer  
 <400> SEQUENCE: 11  
 gacaacagac gcttgacacg 20

-continued

```

<210> SEQ ID NO 12
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 12
ccgactcgag aattcggccc tgcaggccac aacagtctcg cgagttttta attccaagcg      60
ggggttgccc tccgcttggg acaaaaaccg ggggtgcagcc cttggtac                    108

<210> SEQ ID NO 13
<211> LENGTH: 9599
<212> TYPE: DNA
<213> ORGANISM: Hepatitis C virus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (342)..(9377)

<400> SEQUENCE: 13
gccagccccc tgatgggggc gacactccac catgaatcac tcccctgtga ggaactactg      60
tcttcacgca gaaagcgtct agccatggcg ttagtatgag tgtcgtgcag cctccaggac     120
ccccctccc gggagagcca tagtggtctg cggaaccggt gagtacaccg gaattgccag      180
gacgaccggg tcctttcttg gataaacccg ctcaatgcct ggagatttgg gcgtgcccc     240
gcaagactgc tagccgagta gtgttggtgc gcgaaaggcc ttgtgttact gcctgatagg     300
gtgcttgcca gtgccccggg aggtctcgta gaccgtgcac c atg agc acg aat cct      356
                                     Met Ser Thr Asn Pro
                                     1                    5

aaa cct caa aga aaa acc aaa cgt aac acc aac cgt cgc cca cag gac      404
Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn Arg Arg Pro Gln Asp
                                     10                    15                    20

gtc aag ttc ccg ggt ggc ggt cag atc gtt ggt gga gtt tac ttg ttg      452
Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly Gly Val Tyr Leu Leu
                                     25                    30                    35

ccg cgc agg ggc cct aga ttg ggt gtg cgc gcg acg agg aag act tcc      500
Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala Thr Arg Lys Thr Ser
                                     40                    45                    50

gag cgg tcg caa cct cga ggt aga cgt cag cct atc ccc aag gca cgt      548
Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro Ile Pro Lys Ala Arg
                                     55                    60                    65

cgg ccc gag ggc agg acc tgg gct cag ccc ggg tac cct tgg ccc ctc      596
Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly Tyr Pro Trp Pro Leu
                                     70                    75                    80                    85

tat ggc aat gag ggt tgc ggg tgg gcg gga tgg ctc ctg tct ccc cgt      644
Tyr Gly Asn Glu Gly Cys Gly Trp Ala Gly Trp Leu Leu Ser Pro Arg
                                     90                    95                    100

ggc tct cgg cct agc tgg ggc ccc aca gac ccc cgg cgt agg tcg cgc      692
Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro Arg Arg Arg Ser Arg
                                     105                   110                   115

aat ttg ggt aag gtc atc gat acc ctt acg tgc ggc ttc gcc gac ctc      740
Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys Gly Phe Ala Asp Leu
                                     120                   125                   130

atg ggg tac ata ccg ctc gtc ggc gcc cct ctt gga ggc gct gcc agg      788
Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu Gly Gly Ala Ala Arg
                                     135                   140                   145

gcc ctg gcg cat ggc gtc cgg gtt ctg gaa gac ggc gtg aac tat gca      836
Ala Leu Ala His Gly Val Arg Val Leu Glu Asp Gly Val Asn Tyr Ala
                                     150                   155                   160                   165
    
```

-continued

aca ggg aac ctt cct ggt tgc tct ttc tct atc ttc ctt ctg gcc ctg Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile Phe Leu Leu Ala Leu	884
170 175 180	
ctc tct tgc ctg act gtg ccc gct tca gcc tac caa gtg cgc aat tcc Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr Gln Val Arg Asn Ser	932
185 190 195	
tcg ggg ctt tac cat gtc acc aat gat tgc cct aac tcg agt att gtg Ser Gly Leu Tyr His Val Thr Asn Asp Cys Pro Asn Ser Ser Ile Val	980
200 205 210	
tac gag gcg gcc gat gcc atc ctg cac act ccg ggg tgt gtc cct tgc Tyr Glu Ala Ala Asp Ala Ile Leu His Thr Pro Gly Cys Val Pro Cys	1028
215 220 225	
gtt cgc gag ggt aac gcc tcg agg tgt tgg gtg gcg gtg acc ccc acg Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val Ala Val Thr Pro Thr	1076
230 235 240 245	
gtg gcc acc agg gac gcc aaa ctc ccc aca acg cag ctt cga cgt cat Val Ala Thr Arg Asp Gly Lys Leu Pro Thr Thr Gln Leu Arg Arg His	1124
250 255 260	
atc gat ctg ctt gtc ggg agc gcc acc ctc tgc tcg gcc ctc tac gtg Ile Asp Leu Leu Val Gly Ser Ala Thr Leu Cys Ser Ala Leu Tyr Val	1172
265 270 275	
ggg gac ctg tgc ggg tct gtc ttt ctt gtt ggt caa ctg ttt acc ttc Gly Asp Leu Cys Gly Ser Val Phe Leu Val Gly Gln Leu Phe Thr Phe	1220
280 285 290	
tct ccc agg cgc cac tgg acg acg caa gac tgc aat tgt tct atc tat Ser Pro Arg Arg His Trp Thr Thr Gln Asp Cys Asn Cys Ser Ile Tyr	1268
295 300 305	
ccc gcc cat ata acg ggt cat cgc atg gca tgg gat atg atg atg aac Pro Gly His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn	1316
310 315 320 325	
tgg tcc cct acg gca gcg ttg gtg gta gct cag ctg ctc cgg atc cca Trp Ser Pro Thr Ala Ala Leu Val Val Ala Gln Leu Leu Arg Ile Pro	1364
330 335 340	
caa gcc atc atg gac atg atc gct ggt gct cac tgg gga gtc ctg gcg Gln Ala Ile Met Asp Met Ile Ala Gly Ala His Trp Gly Val Leu Ala	1412
345 350 355	
ggc ata gcg tat ttc tcc atg gtg ggg aac tgg gcg aag gtc ctg gta Gly Ile Ala Tyr Phe Ser Met Val Gly Asn Trp Ala Lys Val Leu Val	1460
360 365 370	
gtg ctg ctg cta ttt gcc gcc gtc gac gcg gaa acc cac gtc acc ggg Val Leu Leu Leu Phe Ala Gly Val Asp Ala Glu Thr His Val Thr Gly	1508
375 380 385	
gga aat gcc gcc cgc acc acg gct ggg ctt gtt ggt ctc ctt aca cca Gly Asn Ala Gly Arg Thr Thr Ala Gly Leu Val Gly Leu Leu Thr Pro	1556
390 395 400 405	
ggc gcc aag cag aac atc caa ctg atc aac acc aac gcc agt tgg cac Gly Ala Lys Gln Asn Ile Gln Leu Ile Asn Thr Asn Gly Ser Trp His	1604
410 415 420	
atc aat agc acg gcc ttg aat tgc aat gaa agc ctt aac acc gcc tgg Ile Asn Ser Thr Ala Leu Asn Cys Asn Glu Ser Leu Asn Thr Gly Trp	1652
425 430 435	
tta gca ggg ctc ttc tat caa cac aaa ttc aac tct tca gcc tgt cct Leu Ala Gly Leu Phe Tyr Gln His Lys Phe Asn Ser Ser Gly Cys Pro	1700
440 445 450	
gag agg ttg gcc agc tgc cga cgc ctt acc gat ttt gcc cag gcc tgg Glu Arg Leu Ala Ser Cys Arg Arg Leu Thr Asp Phe Ala Gln Gly Trp	1748
455 460 465	
ggt cct atc agt tat gcc aac gga agc gcc ctc gac gaa cgc ccc tac Gly Pro Ile Ser Tyr Ala Asn Gly Ser Gly Leu Asp Glu Arg Pro Tyr	1796
470 475 480 485	

-continued

tgc tgg cac tac cct cca aga cct tgt ggc att gtg ccc gca aag agc	1844
Cys Trp His Tyr Pro Pro Arg Pro Cys Gly Ile Val Pro Ala Lys Ser	
490 495 500	
gtg tgt ggc ccg gta tat tgc ttc act ccc agc ccc gtg gtg gtg gga	1892
Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val Gly	
505 510 515	
acg acc gac agg tgg ggc gcg cct acc tac agc tgg ggt gca aat gat	1940
Thr Thr Asp Arg Ser Gly Ala Pro Thr Tyr Ser Trp Gly Ala Asn Asp	
520 525 530	
acg gat gtc ttc gtc ctt aac aac acc agg cca ccg ctg ggc aat tgg	1988
Thr Asp Val Phe Val Leu Asn Asn Thr Arg Pro Pro Leu Gly Asn Trp	
535 540 545	
ttc ggt tgt acc tgg atg aac tca act gga ttc acc aaa gtg tgc gga	2036
Phe Gly Cys Thr Trp Met Asn Ser Thr Gly Phe Thr Lys Val Cys Gly	
550 555 560 565	
gcg ccc cct tgt gtc atc gga ggg gtg ggc aac aac acc ttg ctc tgc	2084
Ala Pro Pro Cys Val Ile Gly Gly Val Gly Asn Asn Thr Leu Leu Cys	
570 575 580	
ccc act gat tgc ttc cgc aaa cat ccg gaa gcc aca tac tct cgg tgc	2132
Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ser Arg Cys	
585 590 595	
ggc tcc ggt ccc tgg att aca ccc agg tgc atg gtc gac tac ccg tat	2180
Gly Ser Gly Pro Trp Ile Thr Pro Arg Cys Met Val Asp Tyr Pro Tyr	
600 605 610	
agg ctt tgg cac tat cct tgt acc atc aat tac acc ata ttc aaa gtc	2228
Arg Leu Trp His Tyr Pro Cys Thr Ile Asn Tyr Thr Ile Phe Lys Val	
615 620 625	
agg atg tac gtg gga ggg gtc gag cac agg ctg gaa gcg gcc tgc aac	2276
Arg Met Tyr Val Gly Gly Val Glu His Arg Leu Glu Ala Ala Cys Asn	
630 635 640 645	
tgg acg cgg ggc gaa cgc tgt gat ctg gaa gac agg gac agg tcc gag	2324
Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp Arg Asp Arg Ser Glu	
650 655 660	
ctc agc ccg ttg ctg ctg tcc acc aca cag tgg cag gtc ctt ccg tgt	2372
Leu Ser Pro Leu Leu Leu Ser Thr Thr Gln Trp Gln Val Leu Pro Cys	
665 670 675	
tct ttc acg acc ctg cca gcc ttg tcc acc ggc ctc atc cac ctc cac	2420
Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr Gly Leu Ile His Leu His	
680 685 690	
cag aac att gtg gac gtg cag tac ttg tac ggg gta ggg tca agc atc	2468
Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly Val Gly Ser Ser Ile	
695 700 705	
gcg tcc tgg gcc att aag tgg gag tac gtc gtt ctc ctg ttc ctt ctg	2516
Ala Ser Trp Ala Ile Lys Trp Glu Tyr Val Val Leu Leu Phe Leu Leu	
710 715 720 725	
ctt gca gac gcg cgc gtc tgc tcc tgc ttg tgg atg atg tta ctc ata	2564
Leu Ala Asp Ala Arg Val Cys Ser Cys Leu Trp Met Met Leu Leu Ile	
730 735 740	
tcc caa gcg gag gcg gct ttg gag aac ctc gta ata ctc aat gca gca	2612
Ser Gln Ala Glu Ala Ala Leu Glu Asn Leu Val Ile Leu Asn Ala Ala	
745 750 755	
tcc ctg gcc ggg acg cac ggt ctt gtg tcc ttc ctc gtg ttc ttc tgc	2660
Ser Leu Ala Gly Thr His Gly Leu Val Ser Phe Leu Val Phe Phe Cys	
760 765 770	
ttt gcg tgg tat ctg aag ggt agg tgg gtg ccc gga gcg gtc tac gcc	2708
Phe Ala Trp Tyr Leu Lys Gly Arg Trp Val Pro Gly Ala Val Tyr Ala	
775 780 785	
ctc tac ggg atg tgg cct ctc ctc ctg ctc ctg ctg gcg ttg cct cag	2756
Leu Tyr Gly Met Trp Pro Leu Leu Leu Leu Leu Leu Ala Leu Pro Gln	

-continued

790	795	800	805	
cgg gca tac gca ctg gac acg gag gtg gcc gcg tcg tgt ggc ggc gtt				2804
Arg Ala Tyr Ala Leu Asp Thr Glu Val Ala Ala Ser Cys Gly Gly Val	810	815	820	
gtt ctt gtc ggg tta atg gcg ctg act ctg tcg cca tat tac aag cgc				2852
Val Leu Val Gly Leu Met Ala Leu Thr Leu Ser Pro Tyr Tyr Lys Arg	825	830	835	
tat atc agc tgg tgc atg tgg tgg ctt cag tat ttt ctg acc aga gta				2900
Tyr Ile Ser Trp Cys Met Trp Trp Leu Gln Tyr Phe Leu Thr Arg Val	840	845	850	
gaa gcg caa ctg cac gtg tgg gtt ccc ccc ctc aac gtc cgg ggg ggg				2948
Glu Ala Gln Leu His Val Trp Val Pro Pro Leu Asn Val Arg Gly Gly	855	860	865	
cgc gat gcc gtc atc tta ctc atg tgt gta gta cac ccg acc ctg gta				2996
Arg Asp Ala Val Ile Leu Leu Met Cys Val Val His Pro Thr Leu Val	870	875	880	885
ttt gac atc acc aaa cta ctc ctg gcc atc ttc gga ccc ctt tgg att				3044
Phe Asp Ile Thr Lys Leu Leu Leu Ala Ile Phe Gly Pro Leu Trp Ile	890	895	900	
ctt caa gcc agt ttg ctt aaa gtc ccc tac ttc gtg cgc gtt caa gcc				3092
Leu Gln Ala Ser Leu Leu Lys Val Pro Tyr Phe Val Arg Val Gln Gly	905	910	915	
ctt ctc cgg atc tgc gcg cta gcg cgg aag ata gcc gga ggt cat tac				3140
Leu Leu Arg Ile Cys Ala Leu Ala Arg Lys Ile Ala Gly Gly His Tyr	920	925	930	
gtg caa atg gcc atc atc aag tta ggg gcg ctt act ggc acc tat gtg				3188
Val Gln Met Ala Ile Ile Lys Leu Gly Ala Leu Thr Gly Thr Tyr Val	935	940	945	
tat aac cat ctc acc cct ctt cga gac tgg gcg cac aac ggc ctg cga				3236
Tyr Asn His Leu Thr Pro Leu Arg Asp Trp Ala His Asn Gly Leu Arg	950	955	960	965
gat ctg gcc gtg gct gtg gaa cca gtc gtc ttc tcc cga atg gag acc				3284
Asp Leu Ala Val Ala Val Glu Pro Val Val Phe Ser Arg Met Glu Thr	970	975	980	
aag ctc atc acg tgg ggg gca gat acc gcc gcg tgc ggt gac atc atc				3332
Lys Leu Ile Thr Trp Gly Ala Asp Thr Ala Ala Cys Gly Asp Ile Ile	985	990	995	
aac ggc ttg ccc gtc tct gcc cgt agg ggc cag gag ata ctg ctt ggg				3380
Asn Gly Leu Pro Val Ser Ala Arg Arg Gly Gln Glu Ile Leu Leu Gly	1000	1005	1010	
cca gcc gac gga atg gtc tcc aag ggg tgg agg ttg ctg gcg ccc atc				3428
Pro Ala Asp Gly Met Val Ser Lys Gly Trp Arg Leu Leu Ala Pro Ile	1015	1020	1025	
acg gcg tac gcc cag cag acg aga gcc ctc cta ggg tgt ata atc acc				3476
Thr Ala Tyr Ala Gln Gln Thr Arg Gly Leu Leu Gly Cys Ile Ile Thr	1030	1035	1040	1045
agc ctg act ggc cgg gac aaa aac caa gtg gag ggt gag gtc cag atc				3524
Ser Leu Thr Gly Arg Asp Lys Asn Gln Val Glu Gly Glu Val Gln Ile	1050	1055	1060	
gtg tca act gct acc caa acc ttc ctg gca acg tgc atc aat ggg gta				3572
Val Ser Thr Ala Thr Gln Thr Phe Leu Ala Thr Cys Ile Asn Gly Val	1065	1070	1075	
tgc tgg act gtc tac cac ggg gcc gga acg agg acc atc gca tca ccc				3620
Cys Trp Thr Val Tyr His Gly Ala Gly Thr Arg Thr Ile Ala Ser Pro	1080	1085	1090	
aag ggt cct gtc atc cag atg tat acc aat gtg gac caa gac ctt gtg				3668
Lys Gly Pro Val Ile Gln Met Tyr Thr Asn Val Asp Gln Asp Leu Val	1095	1100	1105	
ggc tgg ccc gct cct caa ggt tcc cgc tca ttg aca ccc tgt acc tgc				3716



-continued

Gly Trp Pro Ala Pro Gln Gly Ser Arg Ser Leu Thr Pro Cys Thr Cys	
1110 1115 1120 1125	
ggc tcc tcg gac ctt tac ctg gtc acg agg cac gcc gat gtc att ccc	3764
Gly Ser Ser Asp Leu Tyr Leu Val Thr Arg His Ala Asp Val Ile Pro	
1130 1135 1140	
gtg cgc cgg cga ggt gat agc agg ggt agc ctg ctt tcg ccc cgg ccc	3812
Val Arg Arg Arg Gly Asp Ser Arg Gly Ser Leu Leu Ser Pro Arg Pro	
1145 1150 1155	
att tcc tac ttg aaa ggc tcc tcg ggg ggt ccg ctg ttg tgc ccc gcg	3860
Ile Ser Tyr Leu Lys Gly Ser Ser Gly Gly Pro Leu Leu Cys Pro Ala	
1160 1165 1170	
gga cac gcc gtg ggc cta ttc agg gcc gcg gtg tgc acc cgt gga gtg	3908
Gly His Ala Val Gly Leu Phe Arg Ala Ala Val Cys Thr Arg Gly Val	
1175 1180 1185	
gct aaa gcg gtg gac ttt atc cct gtg gag aac cta ggg aca acc atg	3956
Ala Lys Val Ala Val Asp Phe Ile Pro Val Glu Asn Leu Gly Thr Thr Met	
1190 1195 1200 1205	
aga tcc ccg gtg ttc acg gac aac tcc tct cca cca gca gtg ccc cag	4004
Arg Ser Pro Val Phe Thr Asp Asn Ser Ser Pro Pro Ala Val Pro Gln	
1210 1215 1220	
agc ttc cag gtg gcc cac ctg cat gct ccc acc gcc agc ggt aag agc	4052
Ser Phe Gln Val Ala His Leu His Ala Pro Thr Gly Ser Gly Lys Ser	
1225 1230 1235	
acc aag gtc ccg gct gcg tac gca gcc cag gcc tac aag gtg ttg gtg	4100
Thr Lys Val Pro Ala Ala Tyr Ala Ala Gln Gly Tyr Lys Val Leu Val	
1240 1245 1250	
ctc aac ccc tct gtt gct gca acg ctg gcc ttt ggt gct tac atg tcc	4148
Leu Asn Pro Ser Val Ala Ala Thr Leu Gly Phe Gly Ala Tyr Met Ser	
1255 1260 1265	
aag gcc cat ggg gtt gat cct aat atc agg acc ggg gtg aga aca att	4196
Lys Ala His Gly Val Asp Pro Asn Ile Arg Thr Gly Val Arg Thr Ile	
1270 1275 1280 1285	
acc act gcc agc ccc atc acg tac tcc acc tac gcc aag ttc ctt gcc	4244
Thr Thr Gly Ser Pro Ile Thr Tyr Ser Thr Tyr Gly Lys Phe Leu Ala	
1290 1295 1300	
gac gcc ggg tgc tca gga ggt gct tat gac ata ata att tgt gac gag	4292
Asp Gly Gly Cys Ser Gly Gly Ala Tyr Asp Ile Ile Ile Cys Asp Glu	
1305 1310 1315	
tgc cac tcc acg gat gcc aca tcc atc ttg gcc atc gcc act gtc ctt	4340
Cys His Ser Thr Asp Ala Thr Ser Ile Leu Gly Ile Gly Thr Val Leu	
1320 1325 1330	
gac caa gca gag act gcg ggg gcg aga ctg gtt gtg ctc gcc act gct	4388
Asp Gln Ala Glu Thr Ala Gly Ala Arg Leu Val Val Leu Ala Thr Ala	
1335 1340 1345	
acc cct ccg gcc tcc gtc act gtg tcc cat cct aac atc gag gag gtt	4436
Thr Pro Pro Gly Ser Val Thr Val Ser His Pro Asn Ile Glu Glu Val	
1350 1355 1360 1365	
gct ctg tcc acc acc gga gag atc ccc ttt tac gcc aag gct atc ccc	4484
Ala Leu Ser Thr Thr Gly Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro	
1370 1375 1380	
ctc gag gtg atc aag ggg gga aga cat ctc atc ttc tgc cac tca aag	4532
Leu Glu Val Ile Lys Gly Gly Arg His Leu Ile Phe Cys His Ser Lys	
1385 1390 1395	
aag aag tgc gac gag ctc gcc gcg aag ctg gtc gca ttg gcc atc aat	4580
Lys Lys Cys Asp Glu Leu Ala Ala Lys Leu Val Ala Leu Gly Ile Asn	
1400 1405 1410	
gcc gtg gcc tac tac cgc ggt ctt gac gtg tct gtc atc ccg acc agc	4628
Ala Val Ala Tyr Tyr Arg Gly Leu Asp Val Ser Val Ile Pro Thr Ser	
1415 1420 1425	

-continued

ggc gat gtt gtc gtc gtg tcg acc gat gct ctc atg act ggc ttt acc Gly Asp Val Val Val Val Ser Thr Asp Ala Leu Met Thr Gly Phe Thr 1430 1435 1440 1445	4676
ggc gac ttc gac tct gtg ata gac tgc aac acg tgt gtc act cag aca Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr 1450 1455 1460	4724
gtc gat ttc agc ctt gac cct acc ttt acc att gag aca acc acg ctc Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu 1465 1470 1475	4772
ccc cag gat gct gtc tcc agg act caa cgc cgg ggc agg act ggc agg Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg 1480 1485 1490	4820
ggg aag cca ggc atc tat aga ttt gtg gca ccg ggg gag cgc ccc tcc Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro Ser 1495 1500 1505	4868
ggc atg ttc gac tcg tcc gtc ctc tgt gag tgc tat gac gcg ggc tgt Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys 1510 1515 1520 1525	4916
gct tgg tat gag ctc acg ccc gcc gag act aca gtt agg cta cga gcg Ala Trp Tyr Glu Leu Thr Pro Ala Glu Thr Thr Val Arg Leu Arg Ala 1530 1535 1540	4964
tac atg aac acc ccg ggg ctt ccc gtg tgc cag gac cat ctt gaa ttt Tyr Met Asn Thr Pro Gly Leu Pro Val Cys Gln Asp His Leu Glu Phe 1545 1550 1555	5012
tgg gag ggc gtc ttt acg ggc ctc act cat ata gat gcc cac ttt tta Trp Glu Gly Val Phe Thr Gly Leu Thr His Ile Asp Ala His Phe Leu 1560 1565 1570	5060
tcc cag aca aag cag agt ggg gag aac ttt cct tac ctg gta gcg tac Ser Gln Thr Lys Gln Ser Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr 1575 1580 1585	5108
caa gcc acc gtg tgc gct agg gct caa gcc cct ccc cca tcg tgg gac Gln Ala Thr Val Cys Ala Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp 1590 1595 1600 1605	5156
cag atg tgg aag tgt ttg atc cgc ctt aaa ccc acc ctc cat ggg cca Gln Met Trp Lys Cys Leu Ile Arg Leu Lys Pro Thr Leu His Gly Pro 1610 1615 1620	5204
aca ccc ctg cta tac aga ctg ggc gct gtt cag aat gaa gtc acc ctg Thr Pro Leu Leu Tyr Arg Leu Gly Ala Val Gln Asn Glu Val Thr Leu 1625 1630 1635	5252
acg cac cca atc acc aaa tac atc atg aca tgc atg tcg gcc gac ctg Thr His Pro Ile Thr Lys Tyr Ile Met Thr Cys Met Ser Ala Asp Leu 1640 1645 1650	5300
gag gtc gtc acg agc acc tgg gtg ctc gtt ggc ggc gtc ctg gct gct Glu Val Val Thr Ser Thr Trp Val Leu Val Gly Gly Val Leu Ala Ala 1655 1660 1665	5348
ctg gcc gcg tat tgc ctg tca aca ggc tgc gtg gtc ata gtg ggc agg Leu Ala Ala Tyr Cys Leu Ser Thr Gly Cys Val Val Ile Val Gly Arg 1670 1675 1680 1685	5396
atc gtc ttg tcc ggg aag ccg gca att ata cct gac agg gag gtt ctc Ile Val Leu Ser Gly Lys Pro Ala Ile Ile Pro Asp Arg Glu Val Leu 1690 1695 1700	5444
tac cag gag ttc gat gag atg gaa gag tgc tct cag cac tta ccg tac Tyr Gln Glu Phe Asp Glu Met Glu Glu Cys Ser Gln His Leu Pro Tyr 1705 1710 1715	5492
atc gag caa ggg atg atg ctc gct gag cag ttc aag cag aag gcc ctc Ile Glu Gln Gly Met Met Leu Ala Glu Gln Phe Lys Gln Lys Ala Leu 1720 1725 1730	5540
ggc ctc ctg cag acc gcg tcc cgc cat gca gag gtt atc acc cct gct Gly Leu Leu Gln Thr Ala Ser Arg His Ala Glu Val Ile Thr Pro Ala 1735 1740 1745	5588

-continued

---

gtc cag acc aac tgg cag aaa ctc gag gtc ttt tgg gcg aag cac atg	5636
Val Gln Thr Asn Trp Gln Lys Leu Glu Val Phe Trp Ala Lys His Met	
1750 1755 1760 1765	
tgg aat ttc atc agt ggg ata caa tac ttg gcg gcc ctg tca acg ctg	5684
Trp Asn Phe Ile Ser Gly Ile Gln Tyr Leu Ala Gly Leu Ser Thr Leu	
1770 1775 1780	
cct ggt aac ccc gcc att gct tca ttg atg gct ttt aca gct gcc gtc	5732
Pro Gly Asn Pro Ala Ile Ala Ser Leu Met Ala Phe Thr Ala Ala Val	
1785 1790 1795	
acc agc cca cta acc act ggc caa acc ctc ctc ttc aac ata ttg ggg	5780
Thr Ser Pro Leu Thr Thr Gly Gln Thr Leu Leu Phe Asn Ile Leu Gly	
1800 1805 1810	
ggg tgg gtg gct gcc cag ctc gcc gcc ccc ggt gcc gct act gcc ttt	5828
Gly Trp Val Ala Ala Gln Leu Ala Ala Pro Gly Ala Ala Thr Ala Phe	
1815 1820 1825	
gtg ggt gct ggc cta gct ggc gcc gcc atc ggc agc gtt gga ctg ggg	5876
Val Gly Ala Gly Leu Ala Gly Ala Ala Ile Gly Ser Val Gly Leu Gly	
1830 1835 1840 1845	
aag gtc ctc gtg gac att ctt gca ggg tat ggc gcg gcc gtg gcg gga	5924
Lys Val Leu Val Asp Ile Leu Ala Gly Tyr Gly Val Ala Gly	
1850 1855 1860	
gct ctt gta gca ttc aag atc atg agc ggt gag gtc ccc tcc acg gag	5972
Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro Ser Thr Glu	
1865 1870 1875	
gac ctg gtc aat ctg ctg ccc gcc atc ctc tcg cct gga gcc ctt gta	6020
Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly Ala Leu Val	
1880 1885 1890	
gtc ggt gtg gtc tgc gca gca ata ctg cgc cgg cac gtt gcc ccg gcc	6068
Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val Gly Pro Gly	
1895 1900 1905	
gag ggg gca gtg caa tgg atg aac cgg cta ata gcc ttc gcc tcc cgg	6116
Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile Ala Phe Ala Ser Arg	
1910 1915 1920 1925	
ggg aac cat gtt tcc ccc acg cac tac gtg ccg gag agc gat gca gcc	6164
Gly Asn His Val Ser Pro Thr His Tyr Val Pro Glu Ser Asp Ala Ala	
1930 1935 1940	
gcc gcg gtc act gcc ata ctc agc agc ctc act gta acc cag ctc ctg	6212
Ala Arg Val Thr Ala Ile Leu Ser Ser Leu Thr Val Thr Gln Leu Leu	
1945 1950 1955	
agg cga ctg cat cag tgg ata agc tcg gag tgt acc act cca tgc tcc	6260
Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser	
1960 1965 1970	
ggt tcc tgg cta agg gac atc tgg gac tgg ata tgc gag gtg ctg agc	6308
Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile Cys Glu Val Leu Ser	
1975 1980 1985	
gac ttt aag acc tgg ctg aaa gcc aag ctc atg cca caa ctg cct ggg	6356
Asp Phe Lys Thr Trp Leu Lys Ala Lys Leu Met Pro Gln Leu Pro Gly	
1990 1995 2000 2005	
att ccc ttt gtg tcc tgc cag cgc ggg tat agg ggg gtc tgg cga gga	6404
Ile Pro Phe Val Ser Cys Gln Arg Gly Tyr Arg Gly Val Trp Arg Gly	
2010 2015 2020	
gac gcc att atg cac act cgc tgc cac tgt gga gct gag atc act gga	6452
Asp Gly Ile Met His Thr Arg Cys His Cys Gly Ala Glu Ile Thr Gly	
2025 2030 2035	
cat gtc aaa aac ggg acg atg agg atc gtc ggt cct agg acc tgc agg	6500
His Val Lys Asn Gly Thr Met Arg Ile Val Gly Pro Arg Thr Cys Arg	
2040 2045 2050	
aac atg tgg agt ggg acg ttc ccc att aac gcc tac acc acg gcc ccc	6548
Asn Met Trp Ser Gly Thr Phe Pro Ile Asn Ala Tyr Thr Thr Gly Pro	

-continued

2055	2060	2065	
tgt act ccc ctt cct gcg cgg aac tat aag ttc gcg ctg tgg agg gtg Cys Thr Pro Leu Pro Ala Pro Asn Tyr Lys Phe Ala Leu Trp Arg Val 2070 2075 2080 2085			6596
tct gca gag gaa tac gtg gag ata agg cgg gtg ggg gac ttc cac tac Ser Ala Glu Glu Tyr Val Glu Ile Arg Arg Val Gly Asp Phe His Tyr 2090 2095 2100			6644
gta tcg ggt atg act act gac aat ctt aaa tgc ccg tgc cag atc cca Val Ser Gly Met Thr Thr Asp Asn Leu Lys Cys Pro Cys Gln Ile Pro 2105 2110 2115			6692
tcg ccc gaa ttt ttc aca gaa ttg gac ggg gtg cgc cta cac agg ttt Ser Pro Glu Phe Phe Thr Glu Leu Asp Gly Val Arg Leu His Arg Phe 2120 2125 2130			6740
gcg ccc cct tgc aag ccc ttg ctg cgg gag gag gta tca ttc aga gta Ala Pro Pro Cys Lys Pro Leu Leu Arg Glu Glu Val Ser Phe Arg Val 2135 2140 2145			6788
gga ctc cac gag tac ccg gtg ggg tcg caa tta cct tgc gag ccc gaa Gly Leu His Glu Tyr Pro Val Gly Ser Gln Leu Pro Cys Glu Pro Glu 2150 2155 2160 2165			6836
ccg gac gta gcc gtg ttg acg tcc atg ctc act gat ccc tcc cat ata Pro Asp Val Ala Val Leu Thr Ser Met Leu Thr Asp Pro Ser His Ile 2170 2175 2180			6884
aca gca gag gcg gcc ggg aga agg ttg gcg aga ggg tca ccc cct tct Thr Ala Glu Ala Ala Gly Arg Arg Leu Ala Arg Gly Ser Pro Pro Ser 2185 2190 2195			6932
atg gcc agc tcc tcg gct agc cag ctg tcc gct cca tct ctc aag gca Met Ala Ser Ser Ser Ala Ser Gln Leu Ser Ala Pro Ser Leu Lys Ala 2200 2205 2210			6980
act tgc acc gcc aac cat gac tcc cct gac gcc gag ctc ata gag gct Thr Cys Thr Ala Asn His Asp Ser Pro Asp Ala Glu Leu Ile Glu Ala 2215 2220 2225			7028
aac ctc ctg tgg agg cag gag atg ggc ggc aac atc acc agg gtt gag Asn Leu Leu Trp Arg Gln Glu Met Gly Gly Asn Ile Thr Arg Val Glu 2230 2235 2240 2245			7076
tca gag aac aaa gtg gtg att ctg gac tcc ttc gat ccg ctt gtg gca Ser Glu Asn Lys Val Val Ile Leu Asp Ser Phe Asp Pro Leu Val Ala 2250 2255 2260			7124
gag gag gat gag cgg gag gtc tcc gta cct gca gaa att ctg cgg aag Glu Glu Asp Glu Arg Glu Val Ser Val Pro Ala Glu Ile Leu Arg Lys 2265 2270 2275			7172
tct cgg aga ttc gcc cgg gcc ctg ccc gtc tgg gcg cgg ccg gac tac Ser Arg Arg Phe Ala Arg Ala Leu Pro Val Trp Ala Arg Pro Asp Tyr 2280 2285 2290			7220
aac ccc ccg cta gta gag acg tgg aaa aag cct gac tac gaa cca cct Asn Pro Pro Leu Val Glu Thr Trp Lys Lys Pro Asp Tyr Glu Pro Pro 2295 2300 2305			7268
gtg gtc cat ggc tgc ccg cta cca cct cca cgg tcc cct cct gtg cct Val Val His Gly Cys Pro Leu Pro Pro Pro Arg Ser Pro Pro Val Pro 2310 2315 2320 2325			7316
ccg cct cgg aaa aag cgt acg gtg gtc ctc acc gaa tca acc cta tct Pro Pro Arg Lys Lys Arg Thr Val Val Leu Thr Glu Ser Thr Leu Ser 2330 2335 2340			7364
act gcc ttg gcc gag ctt gcc acc aaa agt ttt ggc agc tcc tca act Thr Ala Leu Ala Glu Leu Ala Thr Lys Ser Phe Gly Ser Ser Ser Thr 2345 2350 2355			7412
tcc gcc att acg ggc gac aat acg aca aca tcc tct gag ccc gcc cct Ser Gly Ile Thr Gly Asp Asn Thr Thr Thr Ser Ser Glu Pro Ala Pro 2360 2365 2370			7460
tct gcc tgc ccc ccc gac tcc gac gtt gag tcc tat tct tcc atg ccc			7508

-continued

Ser Gly Cys Pro Pro Asp Ser Asp Val Glu Ser Tyr Ser Ser Met Pro 2375 2380 2385	
ccc ctg gag ggg gag cct ggg gat ccg gat ctc agc gac ggg tca tgg Pro Leu Glu Gly Glu Pro Gly Asp Pro Asp Leu Ser Asp Gly Ser Trp 2390 2395 2400 2405	7556
tcg acg gtc agt agt ggg gcc gac acg gaa gat gtc gtg tgc tgc tca Ser Thr Val Ser Ser Gly Ala Asp Thr Glu Asp Val Val Cys Cys Ser 2410 2415 2420	7604
atg tct tat tcc tgg aca ggc gca ctc gtc acc ccg tgc gct gcg gaa Met Ser Tyr Ser Trp Thr Gly Ala Leu Val Thr Pro Cys Ala Ala Glu 2425 2430 2435	7652
gaa caa aaa ctg ccc atc aac gca ctg agc aac tcg ttg cta cgc cat Glu Gln Lys Leu Pro Ile Asn Ala Leu Ser Asn Ser Leu Leu Arg His 2440 2445 2450	7700
cac aat ctg gtg tat tcc acc act tca cgc agt gct tgc caa agg cag His Asn Leu Val Tyr Ser Thr Thr Ser Arg Ser Ala Cys Gln Arg Gln 2455 2460 2465	7748
aag aaa gtc aca ttt gac aga ctg caa gtt ctg gac agc cat tac cag Lys Lys Val Thr Phe Asp Arg Leu Gln Val Leu Asp Ser His Tyr Gln 2470 2475 2480 2485	7796
gac gtg ctc aag gag gtc aaa gca gcg gcg tca aaa gtg aag gct aac Asp Val Leu Lys Glu Val Lys Ala Ala Ala Ser Lys Val Lys Ala Asn 2490 2495 2500	7844
ttg cta tcc gta gag gaa gct tgc agc ctg acg ccc cca cat tca gcc Leu Leu Ser Val Glu Glu Ala Cys Ser Leu Thr Pro Pro His Ser Ala 2505 2510 2515	7892
aaa tcc aag ttt ggc tat ggg gca aaa gac gtc cgt tgc cat gcc aga Lys Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val Arg Cys His Ala Arg 2520 2525 2530	7940
aag gcc gta gcc cac atc aac tcc gtg tgg aaa gac ctt ctg gaa gac Lys Ala Val Ala His Ile Asn Ser Val Trp Lys Asp Leu Leu Glu Asp 2535 2540 2545	7988
agt gta aca cca ata gac act acc atc atg gcc aag aac gag gtt ttc Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn Glu Val Phe 2550 2555 2560 2565	8036
tgc gtt cag cct gag aag ggg ggt cgt aag cca gct cgt ctc atc gtg Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala Arg Leu Ile Val 2570 2575 2580	8084
ttc ccc gac ctg ggc gtg cgc gtg tgc gag aag atg gcc ctg tac gac Phe Pro Asp Leu Gly Val Arg Val Cys Glu Lys Met Ala Leu Tyr Asp 2585 2590 2595	8132
gtg gtt agc aag ctc ccc ctg gcc gtg atg gga agc tcc tac gga ttc Val Val Ser Lys Leu Pro Leu Ala Val Met Gly Ser Ser Tyr Gly Phe 2600 2605 2610	8180
caa tac tca cca gga cag cgg gtt gaa ttc ctc gtg caa gcg tgg aag Gln Tyr Ser Pro Gly Gln Arg Val Glu Phe Leu Val Gln Ala Trp Lys 2615 2620 2625	8228
tcc aag aag acc ccg atg ggg ttc tcg tat gat acc cgc tgt ttt gac Ser Lys Lys Thr Pro Met Gly Phe Ser Tyr Asp Thr Arg Cys Phe Asp 2630 2635 2640 2645	8276
tcc aca gtc act gag agc gac atc cgt acg gag gag gca att tac caa Ser Thr Val Thr Ser Ser Asp Ile Arg Thr Glu Glu Ala Ile Tyr Gln 2650 2655 2660	8324
tgt tgt gac ctg gac ccc caa gcc cgc gtg gcc atc aag tcc ctc act Cys Cys Asp Leu Asp Pro Gln Ala Arg Val Ala Ile Lys Ser Leu Thr 2665 2670 2675	8372
gag agg ctt tat gtt ggg ggc cct ctt acc aat tca agg ggg gaa aac Glu Arg Leu Tyr Val Gly Gly Pro Leu Thr Asn Ser Arg Gly Glu Asn 2680 2685 2690	8420

-continued

tgc ggc tac cgc agg tgc cgc gcg agc ggc gta ctg aca act agc tgt Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val Leu Thr Thr Ser Cys 2695 2700 2705	8468
ggt aac acc ctc act tgc tac atc aag gcc cgg gca gcc tgt cga gcc Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala Arg Ala Ala Cys Arg Ala 2710 2715 2720 2725	8516
gca ggg ctc cag gac tgc acc atg ctc gtg tgt ggc gac gac tta gtc Ala Gly Leu Gln Asp Cys Thr Met Leu Val Cys Gly Asp Asp Leu Val 2730 2735 2740	8564
gtt atc tgt gaa agt gcg ggg gtc cag gag gac gcg gcg agc ctg aga Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp Ala Ala Ser Leu Arg 2745 2750 2755	8612
gcc ttc acg gag gct atg acc agg tac tcc gcc ccc ccc ggg gac ccc Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro Pro Gly Asp Pro 2760 2765 2770	8660
cca caa cca gaa tac gac ttg gag ctt ata aca tca tgc tcc tcc aac Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser Cys Ser Ser Asn 2775 2780 2785	8708
gtg tca gtc gcc cac gac ggc gct gga aag agg gtc tac tac ctt acc Val Ser Val Ala His Asp Gly Ala Gly Lys Arg Val Tyr Tyr Leu Thr 2790 2795 2800 2805	8756
cgt gac cct aca acc ccc ctc gcg aga gcc gcg tgg gag aca gca aga Arg Asp Pro Thr Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg 2810 2815 2820	8804
cac act cca gtc aat tcc tgg cta ggc aac ata atc atg ttt gcc ccc His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile Ile Met Phe Ala Pro 2825 2830 2835	8852
aca ctg tgg gcg agg atg ata ctg atg acc cat ttc ttt agc gtc ctc Thr Leu Trp Ala Arg Met Ile Leu Met Thr His Phe Phe Ser Val Leu 2840 2845 2850	8900
ata gcc agg gat cag ctt gaa cag gct ctt aac tgt gag atc tac gga Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn Cys Glu Ile Tyr Gly 2855 2860 2865	8948
gcc tgc tac tcc ata gaa cca ctg gat cta cct cca atc att caa aga Ala Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro Pro Ile Ile Gln Arg 2870 2875 2880 2885	8996
ctc cat ggc ctc agc gca ttt tca ctc cac agt tac tct cca ggt gaa Leu His Gly Leu Ser Ala Phe Ser Leu His Ser Tyr Ser Pro Gly Glu 2890 2895 2900	9044
atc aat agg gtg gcc gca tgc ctc aga aaa ctt ggg gtc ccg ccc ttg Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu Gly Val Pro Pro Leu 2905 2910 2915	9092
cga gct tgg aga cac cgg gcc cgg agc gtc cgc gct agg ctt ctg tcc Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg Ala Arg Leu Leu Ser 2920 2925 2930	9140
aga gga ggc agg gct gcc ata tgt ggc aag tac ctc ttc aac tgg gca Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr Leu Phe Asn Trp Ala 2935 2940 2945	9188
gta aga aca aag ctc aaa ctc act cca ata gcg gcc gct ggc cgg ctg Val Arg Thr Lys Leu Lys Leu Thr Pro Ile Ala Ala Ala Gly Arg Leu 2950 2955 2960 2965	9236
gac ttg tcc ggt tgg ttc acg gct ggc tac agc ggg gga gac att tat Asp Leu Ser Gly Trp Phe Thr Ala Gly Tyr Ser Gly Gly Asp Ile Tyr 2970 2975 2980	9284
cac agc gtg tct cat gcc cgg ccc cgc tgg ttc tgg ttt tgc cta ctc His Ser Val Ser His Ala Arg Pro Arg Trp Phe Trp Phe Cys Leu Leu 2985 2990 2995	9332
ctg ctc gct gca ggg gta ggc atc tac ctc ctc ccc aac cga tga Leu Leu Ala Ala Gly Val Gly Ile Tyr Leu Leu Pro Asn Arg 3000 3005 3010	9377

-continued

agggtggggg aaacactcgc gcctettaag ccatttcctg tttttttttt tttttttttt 9437  
 ttttttttct tttttttttt ctttcctttc cttctttttt tccttttttt ttccctttctt 9497  
 taatggtggc tccatcttag ccctagtcac ggctagctgt gaaaggtccg tgagccgcat 9557  
 gactgcagag agtgctgata ctggcctctc tgcagatcat gt 9599

<210> SEQ ID NO 14  
 <211> LENGTH: 3011  
 <212> TYPE: PRT  
 <213> ORGANISM: Hepatitis C virus

<400> SEQUENCE: 14

Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn  
 1 5 10 15  
 Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly  
 20 25 30  
 Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala  
 35 40 45  
 Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro  
 50 55 60  
 Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly  
 65 70 75 80  
 Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly Cys Gly Trp Ala Gly Trp  
 85 90 95  
 Leu Leu Ser Pro Arg Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro  
 100 105 110  
 Arg Arg Arg Ser Arg Asn Leu Gly Lys Val Ile Asp Thr Leu Thr Cys  
 115 120 125  
 Gly Phe Ala Asp Leu Met Gly Tyr Ile Pro Leu Val Gly Ala Pro Leu  
 130 135 140  
 Gly Gly Ala Ala Arg Ala Leu Ala His Gly Val Arg Val Leu Glu Asp  
 145 150 155 160  
 Gly Val Asn Tyr Ala Thr Gly Asn Leu Pro Gly Cys Ser Phe Ser Ile  
 165 170 175  
 Phe Leu Leu Ala Leu Leu Ser Cys Leu Thr Val Pro Ala Ser Ala Tyr  
 180 185 190  
 Gln Val Arg Asn Ser Ser Gly Leu Tyr His Val Thr Asn Asp Cys Pro  
 195 200 205  
 Asn Ser Ser Ile Val Tyr Glu Ala Ala Asp Ala Ile Leu His Thr Pro  
 210 215 220  
 Gly Cys Val Pro Cys Val Arg Glu Gly Asn Ala Ser Arg Cys Trp Val  
 225 230 235 240  
 Ala Val Thr Pro Thr Val Ala Thr Arg Asp Gly Lys Leu Pro Thr Thr  
 245 250 255  
 Gln Leu Arg Arg His Ile Asp Leu Leu Val Gly Ser Ala Thr Leu Cys  
 260 265 270  
 Ser Ala Leu Tyr Val Gly Asp Leu Cys Gly Ser Val Phe Leu Val Gly  
 275 280 285  
 Gln Leu Phe Thr Phe Ser Pro Arg Arg His Trp Thr Thr Gln Asp Cys  
 290 295 300  
 Asn Cys Ser Ile Tyr Pro Gly His Ile Thr Gly His Arg Met Ala Trp  
 305 310 315 320  
 Asp Met Met Met Asn Trp Ser Pro Thr Ala Ala Leu Val Val Ala Gln  
 325 330 335

-continued

---

Leu Leu Arg Ile Pro Gln Ala Ile Met Asp Met Ile Ala Gly Ala His  
 340 345 350

Trp Gly Val Leu Ala Gly Ile Ala Tyr Phe Ser Met Val Gly Asn Trp  
 355 360 365

Ala Lys Val Leu Val Val Leu Leu Leu Phe Ala Gly Val Asp Ala Glu  
 370 375 380

Thr His Val Thr Gly Gly Asn Ala Gly Arg Thr Thr Ala Gly Leu Val  
 385 390 395 400

Gly Leu Leu Thr Pro Gly Ala Lys Gln Asn Ile Gln Leu Ile Asn Thr  
 405 410 415

Asn Gly Ser Trp His Ile Asn Ser Thr Ala Leu Asn Cys Asn Glu Ser  
 420 425 430

Leu Asn Thr Gly Trp Leu Ala Gly Leu Phe Tyr Gln His Lys Phe Asn  
 435 440 445

Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys Arg Arg Leu Thr Asp  
 450 455 460

Phe Ala Gln Gly Trp Gly Pro Ile Ser Tyr Ala Asn Gly Ser Gly Leu  
 465 470 475 480

Asp Glu Arg Pro Tyr Cys Trp His Tyr Pro Arg Pro Cys Gly Ile  
 485 490 495

Val Pro Ala Lys Ser Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser  
 500 505 510

Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr Tyr Ser  
 515 520 525

Trp Gly Ala Asn Asp Thr Asp Val Phe Val Leu Asn Asn Thr Arg Pro  
 530 535 540

Pro Leu Gly Asn Trp Phe Gly Cys Thr Trp Met Asn Ser Thr Gly Phe  
 545 550 555 560

Thr Lys Val Cys Gly Ala Pro Pro Cys Val Ile Gly Gly Val Gly Asn  
 565 570 575

Asn Thr Leu Leu Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala  
 580 585 590

Thr Tyr Ser Arg Cys Gly Ser Gly Pro Trp Ile Thr Pro Arg Cys Met  
 595 600 605

Val Asp Tyr Pro Tyr Arg Leu Trp His Tyr Pro Cys Thr Ile Asn Tyr  
 610 615 620

Thr Ile Phe Lys Val Arg Met Tyr Val Gly Gly Val Glu His Arg Leu  
 625 630 635 640

Glu Ala Ala Cys Asn Trp Thr Arg Gly Glu Arg Cys Asp Leu Glu Asp  
 645 650 655

Arg Asp Arg Ser Glu Leu Ser Pro Leu Leu Leu Ser Thr Thr Gln Trp  
 660 665 670

Gln Val Leu Pro Cys Ser Phe Thr Thr Leu Pro Ala Leu Ser Thr Gly  
 675 680 685

Leu Ile His Leu His Gln Asn Ile Val Asp Val Gln Tyr Leu Tyr Gly  
 690 695 700

Val Gly Ser Ser Ile Ala Ser Trp Ala Ile Lys Trp Glu Tyr Val Val  
 705 710 715 720

Leu Leu Phe Leu Leu Leu Ala Asp Ala Arg Val Cys Ser Cys Leu Trp  
 725 730 735

Met Met Leu Leu Ile Ser Gln Ala Glu Ala Ala Leu Glu Asn Leu Val  
 740 745 750



-continued

---

Ile Leu Asn Ala Ala Ser Leu Ala Gly Thr His Gly Leu Val Ser Phe  
755 760 765

Leu Val Phe Phe Cys Phe Ala Trp Tyr Leu Lys Gly Arg Trp Val Pro  
770 775 780

Gly Ala Val Tyr Ala Leu Tyr Gly Met Trp Pro Leu Leu Leu Leu Leu  
785 790 795 800

Leu Ala Leu Pro Gln Arg Ala Tyr Ala Leu Asp Thr Glu Val Ala Ala  
805 810 815

Ser Cys Gly Gly Val Val Leu Val Gly Leu Met Ala Leu Thr Leu Ser  
820 825 830

Pro Tyr Tyr Lys Arg Tyr Ile Ser Trp Cys Met Trp Trp Leu Gln Tyr  
835 840 845

Phe Leu Thr Arg Val Glu Ala Gln Leu His Val Trp Val Pro Pro Leu  
850 855 860

Asn Val Arg Gly Gly Arg Asp Ala Val Ile Leu Leu Met Cys Val Val  
865 870 875 880

His Pro Thr Leu Val Phe Asp Ile Thr Lys Leu Leu Leu Ala Ile Phe  
885 890 895

Gly Pro Leu Trp Ile Leu Gln Ala Ser Leu Leu Lys Val Pro Tyr Phe  
900 905 910

Val Arg Val Gln Gly Leu Leu Arg Ile Cys Ala Leu Ala Arg Lys Ile  
915 920 925

Ala Gly Gly His Tyr Val Gln Met Ala Ile Ile Lys Leu Gly Ala Leu  
930 935 940

Thr Gly Thr Tyr Val Tyr Asn His Leu Thr Pro Leu Arg Asp Trp Ala  
945 950 955 960

His Asn Gly Leu Arg Asp Leu Ala Val Ala Val Glu Pro Val Val Phe  
965 970 975

Ser Arg Met Glu Thr Lys Leu Ile Thr Trp Gly Ala Asp Thr Ala Ala  
980 985 990

Cys Gly Asp Ile Ile Asn Gly Leu Pro Val Ser Ala Arg Arg Gly Gln  
995 1000 1005

Glu Ile Leu Leu Gly Pro Ala Asp Gly Met Val Ser Lys Gly Trp Arg  
1010 1015 1020

Leu Leu Ala Pro Ile Thr Ala Tyr Ala Gln Gln Thr Arg Gly Leu Leu  
1025 1030 1035 1040

Gly Cys Ile Ile Thr Ser Leu Thr Gly Arg Asp Lys Asn Gln Val Glu  
1045 1050 1055

Gly Glu Val Gln Ile Val Ser Thr Ala Thr Gln Thr Phe Leu Ala Thr  
1060 1065 1070

Cys Ile Asn Gly Val Cys Trp Thr Val Tyr His Gly Ala Gly Thr Arg  
1075 1080 1085

Thr Ile Ala Ser Pro Lys Gly Pro Val Ile Gln Met Tyr Thr Asn Val  
1090 1095 1100

Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly Ser Arg Ser Leu  
1105 1110 1115 1120

Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr Leu Val Thr Arg His  
1125 1130 1135

Ala Asp Val Ile Pro Val Arg Arg Arg Gly Asp Ser Arg Gly Ser Leu  
1140 1145 1150

Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly Ser Ser Gly Gly Pro  
1155 1160 1165

Leu Leu Cys Pro Ala Gly His Ala Val Gly Leu Phe Arg Ala Ala Val

-continued

1170		1175		1180	
Cys Thr Arg Gly Val Ala Lys Ala Val Asp Phe Ile Pro Val Glu Asn					
1185		1190		1195	1200
Leu Gly Thr Thr Met Arg Ser Pro Val Phe Thr Asp Asn Ser Ser Pro		1205		1210	1215
Pro Ala Val Pro Gln Ser Phe Gln Val Ala His Leu His Ala Pro Thr		1220		1225	1230
Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala Tyr Ala Ala Gln Gly		1235		1240	1245
Tyr Lys Val Leu Val Leu Asn Pro Ser Val Ala Ala Thr Leu Gly Phe		1250		1255	1260
Gly Ala Tyr Met Ser Lys Ala His Gly Val Asp Pro Asn Ile Arg Thr		1265		1270	1275
Gly Val Arg Thr Ile Thr Thr Gly Ser Pro Ile Thr Tyr Ser Thr Tyr		1285		1290	1295
Gly Lys Phe Leu Ala Asp Gly Gly Cys Ser Gly Gly Ala Tyr Asp Ile		1300		1305	1310
Ile Ile Cys Asp Glu Cys His Ser Thr Asp Ala Thr Ser Ile Leu Gly		1315		1320	1325
Ile Gly Thr Val Leu Asp Gln Ala Glu Thr Ala Gly Ala Arg Leu Val		1330		1335	1340
Val Leu Ala Thr Ala Thr Pro Pro Gly Ser Val Thr Val Ser His Pro		1345		1350	1355
Asn Ile Glu Glu Val Ala Leu Ser Thr Thr Gly Glu Ile Pro Phe Tyr		1365		1370	1375
Gly Lys Ala Ile Pro Leu Glu Val Ile Lys Gly Gly Arg His Leu Ile		1380		1385	1390
Phe Cys His Ser Lys Lys Lys Cys Asp Glu Leu Ala Ala Lys Leu Val		1395		1400	1405
Ala Leu Gly Ile Asn Ala Val Ala Tyr Tyr Arg Gly Leu Asp Val Ser		1410		1415	1420
Val Ile Pro Thr Ser Gly Asp Val Val Val Val Ser Thr Asp Ala Leu		1425		1430	1435
Met Thr Gly Phe Thr Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr		1445		1450	1455
Cys Val Thr Gln Thr Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile		1460		1465	1470
Glu Thr Thr Thr Leu Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg		1475		1480	1485
Gly Arg Thr Gly Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro		1490		1495	1500
Gly Glu Arg Pro Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys		1505		1510	1515
Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu Thr Pro Ala Glu Thr Thr		1525		1530	1535
Val Arg Leu Arg Ala Tyr Met Asn Thr Pro Gly Leu Pro Val Cys Gln		1540		1545	1550
Asp His Leu Glu Phe Trp Glu Gly Val Phe Thr Gly Leu Thr His Ile		1555		1560	1565
Asp Ala His Phe Leu Ser Gln Thr Lys Gln Ser Gly Glu Asn Phe Pro		1570		1575	1580

-continued

---

Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys Ala Arg Ala Gln Ala Pro  
 1585 1590 1595 1600  
 Pro Pro Ser Trp Asp Gln Met Trp Lys Cys Leu Ile Arg Leu Lys Pro  
 1605 1610 1615  
 Thr Leu His Gly Pro Thr Pro Leu Leu Tyr Arg Leu Gly Ala Val Gln  
 1620 1625 1630  
 Asn Glu Val Thr Leu Thr His Pro Ile Thr Lys Tyr Ile Met Thr Cys  
 1635 1640 1645  
 Met Ser Ala Asp Leu Glu Val Val Thr Ser Thr Trp Val Leu Val Gly  
 1650 1655 1660  
 Gly Val Leu Ala Ala Leu Ala Ala Tyr Cys Leu Ser Thr Gly Cys Val  
 1665 1670 1675 1680  
 Val Ile Val Gly Arg Ile Val Leu Ser Gly Lys Pro Ala Ile Ile Pro  
 1685 1690 1695  
 Asp Arg Glu Val Leu Tyr Gln Glu Phe Asp Glu Met Glu Glu Cys Ser  
 1700 1705 1710  
 Gln His Leu Pro Tyr Ile Glu Gln Gly Met Met Leu Ala Glu Gln Phe  
 1715 1720 1725  
 Lys Gln Lys Ala Leu Gly Leu Leu Gln Thr Ala Ser Arg His Ala Glu  
 1730 1735 1740  
 Val Ile Thr Pro Ala Val Gln Thr Asn Trp Gln Lys Leu Glu Val Phe  
 1745 1750 1755 1760  
 Trp Ala Lys His Met Trp Asn Phe Ile Ser Gly Ile Gln Tyr Leu Ala  
 1765 1770 1775  
 Gly Leu Ser Thr Leu Pro Gly Asn Pro Ala Ile Ala Ser Leu Met Ala  
 1780 1785 1790  
 Phe Thr Ala Ala Val Thr Ser Pro Leu Thr Thr Gly Gln Thr Leu Leu  
 1795 1800 1805  
 Phe Asn Ile Leu Gly Gly Trp Val Ala Ala Gln Leu Ala Ala Pro Gly  
 1810 1815 1820  
 Ala Ala Thr Ala Phe Val Gly Ala Gly Leu Ala Gly Ala Ala Ile Gly  
 1825 1830 1835 1840  
 Ser Val Gly Leu Gly Lys Val Leu Val Asp Ile Leu Ala Gly Tyr Gly  
 1845 1850 1855  
 Ala Gly Val Ala Gly Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu  
 1860 1865 1870  
 Val Pro Ser Thr Glu Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser  
 1875 1880 1885  
 Pro Gly Ala Leu Val Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg  
 1890 1895 1900  
 His Val Gly Pro Gly Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile  
 1905 1910 1915 1920  
 Ala Phe Ala Ser Arg Gly Asn His Val Ser Pro Thr His Tyr Val Pro  
 1925 1930 1935  
 Glu Ser Asp Ala Ala Ala Arg Val Thr Ala Ile Leu Ser Ser Leu Thr  
 1940 1945 1950  
 Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys  
 1955 1960 1965  
 Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile  
 1970 1975 1980

-continued

---

Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu Lys Ala Lys Leu Met  
 1985 1990 1995 2000  
 Pro Gln Leu Pro Gly Ile Pro Phe Val Ser Cys Gln Arg Gly Tyr Arg  
 2005 2010 2015  
 Gly Val Trp Arg Gly Asp Gly Ile Met His Thr Arg Cys His Cys Gly  
 2020 2025 2030  
 Ala Glu Ile Thr Gly His Val Lys Asn Gly Thr Met Arg Ile Val Gly  
 2035 2040 2045  
 Pro Arg Thr Cys Arg Asn Met Trp Ser Gly Thr Phe Pro Ile Asn Ala  
 2050 2055 2060  
 Tyr Thr Thr Gly Pro Cys Thr Pro Leu Pro Ala Pro Asn Tyr Lys Phe  
 2065 2070 2075 2080  
 Ala Leu Trp Arg Val Ser Ala Glu Glu Tyr Val Glu Ile Arg Arg Val  
 2085 2090 2095  
 Gly Asp Phe His Tyr Val Ser Gly Met Thr Thr Asp Asn Leu Lys Cys  
 2100 2105 2110  
 Pro Cys Gln Ile Pro Ser Pro Glu Phe Phe Thr Glu Leu Asp Gly Val  
 2115 2120 2125  
 Arg Leu His Arg Phe Ala Pro Pro Cys Lys Pro Leu Leu Arg Glu Glu  
 2130 2135 2140  
 Val Ser Phe Arg Val Gly Leu His Glu Tyr Pro Val Gly Ser Gln Leu  
 2145 2150 2155 2160  
 Pro Cys Glu Pro Glu Pro Asp Val Ala Val Leu Thr Ser Met Leu Thr  
 2165 2170 2175  
 Asp Pro Ser His Ile Thr Ala Glu Ala Ala Gly Arg Arg Leu Ala Arg  
 2180 2185 2190  
 Gly Ser Pro Pro Ser Met Ala Ser Ser Ala Ser Gln Leu Ser Ala  
 2195 2200 2205  
 Pro Ser Leu Lys Ala Thr Cys Thr Ala Asn His Asp Ser Pro Asp Ala  
 2210 2215 2220  
 Glu Leu Ile Glu Ala Asn Leu Leu Trp Arg Gln Glu Met Gly Gly Asn  
 2225 2230 2235 2240  
 Ile Thr Arg Val Glu Ser Glu Asn Lys Val Val Ile Leu Asp Ser Phe  
 2245 2250 2255  
 Asp Pro Leu Val Ala Glu Glu Asp Glu Arg Glu Val Ser Val Pro Ala  
 2260 2265 2270  
 Glu Ile Leu Arg Lys Ser Arg Arg Phe Ala Arg Ala Leu Pro Val Trp  
 2275 2280 2285  
 Ala Arg Pro Asp Tyr Asn Pro Pro Leu Val Glu Thr Trp Lys Lys Pro  
 2290 2295 2300  
 Asp Tyr Glu Pro Pro Val Val His Gly Cys Pro Leu Pro Pro Arg  
 2305 2310 2315 2320  
 Ser Pro Pro Val Pro Pro Arg Lys Lys Arg Thr Val Val Leu Thr  
 2325 2330 2335  
 Glu Ser Thr Leu Ser Thr Ala Leu Ala Glu Leu Ala Thr Lys Ser Phe  
 2340 2345 2350  
 Gly Ser Ser Thr Ser Gly Ile Thr Gly Asp Asn Thr Thr Thr Ser  
 2355 2360 2365  
 Ser Glu Pro Ala Pro Ser Gly Cys Pro Pro Asp Ser Asp Val Glu Ser  
 2370 2375 2380  
 Tyr Ser Ser Met Pro Pro Leu Glu Gly Glu Pro Gly Asp Pro Asp Leu  
 2385 2390 2395 2400

-continued

---

Ser Asp Gly Ser Trp Ser Thr Val Ser Ser Gly Ala Asp Thr Glu Asp  
 2405 2410 2415  
 Val Val Cys Cys Ser Met Ser Tyr Ser Trp Thr Gly Ala Leu Val Thr  
 2420 2425 2430  
 Pro Cys Ala Ala Glu Glu Gln Lys Leu Pro Ile Asn Ala Leu Ser Asn  
 2435 2440 2445  
 Ser Leu Leu Arg His His Asn Leu Val Tyr Ser Thr Thr Ser Arg Ser  
 2450 2455 2460  
 Ala Cys Gln Arg Gln Lys Lys Val Thr Phe Asp Arg Leu Gln Val Leu  
 2465 2470 2475 2480  
 Asp Ser His Tyr Gln Asp Val Leu Lys Glu Val Lys Ala Ala Ala Ser  
 2485 2490 2495  
 Lys Val Lys Ala Asn Leu Leu Ser Val Glu Glu Ala Cys Ser Leu Thr  
 2500 2505 2510  
 Pro Pro His Ser Ala Lys Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val  
 2515 2520 2525  
 Arg Cys His Ala Arg Lys Ala Val Ala His Ile Asn Ser Val Trp Lys  
 2530 2535 2540  
 Asp Leu Leu Glu Asp Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala  
 2545 2550 2555 2560  
 Lys Asn Glu Val Phe Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro  
 2565 2570 2575  
 Ala Arg Leu Ile Val Phe Pro Asp Leu Gly Val Arg Val Cys Glu Lys  
 2580 2585 2590  
 Met Ala Leu Tyr Asp Val Val Ser Lys Leu Pro Leu Ala Val Met Gly  
 2595 2600 2605  
 Ser Ser Tyr Gly Phe Gln Tyr Ser Pro Gly Gln Arg Val Glu Phe Leu  
 2610 2615 2620  
 Val Gln Ala Trp Lys Ser Lys Lys Thr Pro Met Gly Phe Ser Tyr Asp  
 2625 2630 2635 2640  
 Thr Arg Cys Phe Asp Ser Thr Val Thr Glu Ser Asp Ile Arg Thr Glu  
 2645 2650 2655  
 Glu Ala Ile Tyr Gln Cys Cys Asp Leu Asp Pro Gln Ala Arg Val Ala  
 2660 2665 2670  
 Ile Lys Ser Leu Thr Glu Arg Leu Tyr Val Gly Gly Pro Leu Thr Asn  
 2675 2680 2685  
 Ser Arg Gly Glu Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val  
 2690 2695 2700  
 Leu Thr Thr Ser Cys Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala Arg  
 2705 2710 2715 2720  
 Ala Ala Cys Arg Ala Ala Gly Leu Gln Asp Cys Thr Met Leu Val Cys  
 2725 2730 2735  
 Gly Asp Asp Leu Val Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp  
 2740 2745 2750  
 Ala Ala Ser Leu Arg Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala  
 2755 2760 2765  
 Pro Pro Gly Asp Pro Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr  
 2770 2775 2780  
 Ser Cys Ser Ser Asn Val Ser Val Ala His Asp Gly Ala Gly Lys Arg  
 2785 2790 2795 2800  
 Val Tyr Tyr Leu Thr Arg Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala  
 2805 2810 2815

-continued

Trp Glu Thr Ala Arg His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile  
 2820 2825 2830

Ile Met Phe Ala Pro Thr Leu Trp Ala Arg Met Ile Leu Met Thr His  
 2835 2840 2845

Phe Phe Ser Val Leu Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn  
 2850 2855 2860

Cys Glu Ile Tyr Gly Ala Cys Tyr Ser Ile Glu Pro Leu Asp Leu Pro  
 2865 2870 2875 2880

Pro Ile Ile Gln Arg Leu His Gly Leu Ser Ala Phe Ser Leu His Ser  
 2885 2890 2895

Tyr Ser Pro Gly Glu Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu  
 2900 2905 2910

Gly Val Pro Pro Leu Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg  
 2915 2920 2925

Ala Arg Leu Leu Ser Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr  
 2930 2935 2940

Leu Phe Asn Trp Ala Val Arg Thr Lys Leu Lys Leu Thr Pro Ile Ala  
 2945 2950 2955 2960

Ala Ala Gly Arg Leu Asp Leu Ser Gly Trp Phe Thr Ala Gly Tyr Ser  
 2965 2970 2975

Gly Gly Asp Ile Tyr His Ser Val Ser His Ala Arg Pro Arg Trp Phe  
 2980 2985 2990

Trp Phe Cys Leu Leu Leu Ala Ala Gly Val Gly Ile Tyr Leu Leu  
 2995 3000 3005

Pro Asn Arg  
 3010

<210> SEQ ID NO 15  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 15

gaaggagga ggtttgaagg a 21

<210> SEQ ID NO 16  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 16

ccagttccgg gcaagaact 19

<210> SEQ ID NO 17  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 Primer

<400> SEQUENCE: 17

aattcgccc tgcaggccac aacagtc 27

87

What is claimed is:

1. A method of producing a recombinant or chimeric GBV-B virus comprising:
  - introducing into a host cell a recombinant GBV-B or chimeric GBV-B viral genome comprising a 3' terminal sequence of GBV-B, wherein the 3' terminal sequence comprises 50 contiguous nucleotides from SEQ ID NO:1; and
  - culturing said host cell under conditions where the 3' terminal sequence permits production of the virus from said genome.
2. The method of claim 1, wherein said 3' terminal sequence comprises 100 contiguous nucleotides from SEQ ID NO:1.
3. The method of claim 2, wherein said 3' terminal sequence comprises SEQ ID NO:1.
4. The method of claim 1, wherein said host cell is in an animal.
5. The method of claim 1, wherein said genome comprises recombinant RNA.
6. The method of claim 1, wherein said genome is encoded by recombinant DNA.
7. The method of claim 1, further comprising the step of isolating virus from said host cell.
8. The method of claim 7, wherein said virus is purified to homogeneity.
9. A method of producing a GBV-B or chimeric GBV-B virus comprising:
  - obtaining a virus produced by the method of claim 1;
  - introducing the virus into a second host cell; and
  - culturing said host cell under conditions permitting production of virus.
10. A method of producing a virus comprising:
  - introducing into a host cell a recombinant GBV-B or chimeric GBV-B viral genome comprising a 3' terminal sequence of GBV-B, wherein the 3' terminal sequence has at least 70% sequence identity to SEQ ID NO:1; and
  - maintaining said host cell under conditions permitting production of a virus from said genome.
11. The method of claim 10, wherein the host cell is a liver cell.
12. The method of claim 11, wherein the liver cell is in an animal.
13. The method of claim 12, wherein the animal is a primate.
14. The method of claim 13, wherein the primate is a non-human primate.

88

15. The method of claim 13, wherein the primate is a tamarin.
16. The method of claim 11, wherein the 3' terminal sequence has at least 80% sequence identity to SEQ ID NO:1.
17. The method of claim 16, wherein the 3' terminal sequence has at least 90% sequence identity to SEQ ID NO:1.
18. The method of claim 17, wherein the 3' terminal sequence has at least 95% sequence identity to SEQ ID NO:1.
19. The method of claim 18, wherein the 3' terminal sequence is SEQ ID NO:1.
20. A method of producing a recombinant or chimeric GBV-B virus comprising:
  - introducing into a liver cell a recombinant GBV-B or chimeric GBV-B viral genome comprising a 3' terminal sequence of GBV-B, wherein the 3' terminal sequence has at least 70% sequence identity to SEQ ID NO:1; and
  - maintaining said liver cell under conditions where the 3' terminal sequence permits production of the virus from said genome.
21. A method of replicating a recombinant chimeric GBV-B virus comprising:
  - introducing into a liver cell a recombinant GBV-B or chimeric GBV-B viral genome comprising a 3' terminal sequence of GBV-B, wherein the 3' terminal sequence has at least 70% sequence identity to SEQ ID NO:1; and
  - maintaining said liver cell under conditions where the 3' terminal sequence permits replication of the recombinant viral genome.
22. The method of claim 21, wherein the 3' terminal sequence has at least 80% sequence identity to SEQ ID NO:1.
23. The method of claim 22, wherein the 3' terminal sequence has at least 90% sequence identity to SEQ ID NO:1.
24. The method of claim 23, wherein the 3' terminal sequence has at least 95% sequence identity to SEQ ID NO:1.
25. The method of claim 24, wherein the 3' terminal sequence is SEQ ID NO:1.

\* \* \* \* \*